

Dissociative disorders. Severe dissociative responses (in the absence of characteristic acute stress disorder symptoms) may be diagnosed as derealization/depersonalization disorder. If severe amnesia of the trauma persists in the absence of characteristic acute stress disorder symptoms, the diagnosis of dissociative amnesia may be indicated.

Posttraumatic stress disorder. Acute stress disorder is distinguished from PTSD because the symptom pattern in acute stress disorder must occur within 1 month of the traumatic event and resolve within that 1-month period. If the symptoms persist for more than 1 month and meet criteria for PTSD, the diagnosis is changed from acute stress disorder to PTSD.

Obsessive-compulsive disorder. In obsessive-compulsive disorder, there are recurrent intrusive thoughts, but these meet the definition of an obsession. In addition, the intrusive thoughts are not related to an experienced traumatic event, compulsions are usually present, and other symptoms of acute stress disorder are typically absent.

Psychotic disorders. Flashbacks in acute stress disorder must be distinguished from illusions, hallucinations, and other perceptual disturbances that may occur in schizophrenia, other psychotic disorders, depressive or bipolar disorder with psychotic features, a delirium, substance/medication-induced disorders, and psychotic disorders due to another medical condition. Acute stress disorder flashbacks are distinguished from these other perceptual disturbances by being directly related to the traumatic experience and by occurring in the absence of other psychotic or substance-induced features.

Traumatic brain injury. When a brain injury occurs in the context of a traumatic event (e.g., traumatic accident, bomb blast, acceleration/deceleration trauma), symptoms of acute stress disorder may appear. An event causing head trauma may also constitute a psychological traumatic event, and traumatic brain injury (TBI)-related neurocognitive symptoms are not mutually exclusive and may occur concurrently. Symptoms previously termed *postconcussive* (e.g., headaches, dizziness, sensitivity to light or sound, irritability, concentration deficits) can occur in brain-injured and non-brain injured populations, including individuals with acute stress disorder. Because symptoms of acute stress disorder and TBI-related neurocognitive symptoms can overlap, a differential diagnosis between acute stress disorder and neurocognitive disorder symptoms attributable to TBI may be possible based on the presence of symptoms that are distinctive to each presentation. Whereas reexperiencing and avoidance are characteristic of acute stress disorder and not the effects of TBI, persistent disorientation and confusion are more specific to TBI (neurocognitive effects) than to acute stress disorder. Furthermore, differential is aided by the fact that symptoms of acute stress disorder persist for up to only 1 month following trauma exposure.

Adjustment Disorders

Diagnostic Criteria

- A. The development of emotional or behavioral symptoms in response to an identifiable stressor(s) occurring within 3 months of the onset of the stressor(s).
- B. These symptoms or behaviors are clinically significant, as evidenced by one or both of the following:
 1. Marked distress that is out of proportion to the severity or intensity of the stressor, taking into account the external context and the cultural factors that might influence symptom severity and presentation.
 2. Significant impairment in social, occupational, or other important areas of functioning.
- C. The stress-related disturbance does not meet the criteria for another mental disorder and is not merely an exacerbation of a preexisting mental disorder.

- D. The symptoms do not represent normal bereavement.
- E. Once the stressor or its consequences have terminated, the symptoms do not persist for more than an additional 6 months.

Specify whether:

309.0 (F43.21) With depressed mood: Low mood, tearfulness, or feelings of hopelessness are predominant.

309.24 (F43.22) With anxiety: Nervousness, worry, jitteriness, or separation anxiety is predominant.

309.28 (F43.23) With mixed anxiety and depressed mood: A combination of depression and anxiety is predominant.

309.3 (F43.24) With disturbance of conduct: Disturbance of conduct is predominant.

309.4 (F43.25) With mixed disturbance of emotions and conduct: Both emotional symptoms (e.g., depression, anxiety) and a disturbance of conduct are predominant.

309.9 (F43.20) Unspecified: For maladaptive reactions that are not classifiable as one of the specific subtypes of adjustment disorder.

Diagnostic Features

The presence of emotional or behavioral symptoms in response to an identifiable stressor is the essential feature of adjustment disorders (Criterion A). The stressor may be a single event (e.g., a termination of a romantic relationship), or there may be multiple stressors (e.g., marked business difficulties and marital problems). Stressors may be recurrent (e.g., associated with seasonal business crises, unfulfilling sexual relationships) or continuous (e.g., a persistent painful illness with increasing disability, living in a crime-ridden neighborhood). Stressors may affect a single individual, an entire family, or a larger group or community (e.g., a natural disaster). Some stressors may accompany specific developmental events (e.g., going to school, leaving a parental home, reentering a parental home, getting married, becoming a parent, failing to attain occupational goals, retirement).

Adjustment disorders may be diagnosed following the death of a loved one when the intensity, quality, or persistence of grief reactions exceeds what normally might be expected, when cultural, religious, or age-appropriate norms are taken into account. A more specific set of bereavement-related symptoms has been designated *persistent complex bereavement disorder*.

Adjustment disorders are associated with an increased risk of suicide attempts and completed suicide.

Prevalence

Adjustment disorders are common, although prevalence may vary widely as a function of the population studied and the assessment methods used. The percentage of individuals in outpatient mental health treatment with a principal diagnosis of an adjustment disorder ranges from approximately 5% to 20%. In a hospital psychiatric consultation setting, it is often the most common diagnosis, frequently reaching 50%.

Development and Course

By definition, the disturbance in adjustment disorders begins within 3 months of onset of a stressor and lasts no longer than 6 months after the stressor or its consequences have ceased. If the stressor is an acute event (e.g., being fired from a job), the onset of the disturbance is usually immediate (i.e., within a few days) and the duration is relatively brief (i.e., no more than a few months). If the stressor or its consequences persist, the adjustment disorder may also continue to be present and become the persistent form.

Risk and Prognostic Factors

Environmental. Individuals from disadvantaged life circumstances experience a high rate of stressors and may be at increased risk for adjustment disorders.

Culture-Related Diagnostic Issues

The context of the individual's cultural setting should be taken into account in making the clinical judgment of whether the individual's response to the stressor is maladaptive or whether the associated distress is in excess of what would be expected. The nature, meaning, and experience of the stressors and the evaluation of the response to the stressors may vary across cultures.

Functional Consequences of Adjustment Disorders

The subjective distress or impairment in functioning associated with adjustment disorders is frequently manifested as decreased performance at work or school and temporary changes in social relationships. An adjustment disorder may complicate the course of illness in individuals who have a general medical condition (e.g., decreased compliance with the recommended medical regimen; increased length of hospital stay).

Differential Diagnosis

Major depressive disorder. If an individual has symptoms that meet criteria for a major depressive disorder in response to a stressor, the diagnosis of an adjustment disorder is not applicable. The symptom profile of major depressive disorder differentiates it from adjustment disorders.

Posttraumatic stress disorder and acute stress disorder. In adjustment disorders, the stressor can be of any severity rather than of the severity and type required by Criterion A of acute stress disorder and posttraumatic stress disorder (PTSD). In distinguishing adjustment disorders from these two posttraumatic diagnoses, there are both timing and symptom profile considerations. Adjustment disorders can be diagnosed immediately and persist up to 6 months after exposure to the traumatic event, whereas acute stress disorder can only occur between 3 days and 1 month of exposure to the stressor, and PTSD cannot be diagnosed until at least 1 month has passed since the occurrence of the traumatic stressor. The required symptom profile for PTSD and acute stress disorder differentiates them from the adjustment disorders. With regard to symptom profiles, an adjustment disorder may be diagnosed following a traumatic event when an individual exhibits symptoms of either acute stress disorder or PTSD that do not meet or exceed the diagnostic threshold for either disorder. An adjustment disorder should also be diagnosed for individuals who have not been exposed to a traumatic event but who otherwise exhibit the full symptom profile of either acute stress disorder or PTSD.

Personality disorders. With regard to personality disorders, some personality features may be associated with a vulnerability to situational distress that may resemble an adjustment disorder. The lifetime history of personality functioning will help inform the interpretation of distressed behaviors to aid in distinguishing a long-standing personality disorder from an adjustment disorder. In addition to some personality disorders incurring vulnerability to distress, stressors may also exacerbate personality disorder symptoms. In the presence of a personality disorder, if the symptom criteria for an adjustment disorder are met, and the stress-related disturbance exceeds what may be attributable to maladaptive personality disorder symptoms (i.e., Criterion C is met), then the diagnosis of an adjustment disorder should be made.

Psychological factors affecting other medical conditions. In psychological factors affecting other medical conditions, specific psychological entities (e.g., psychological symptoms, behaviors, other factors) exacerbate a medical condition. These psychological factors can precipitate, exacerbate, or put an individual at risk for medical illness, or they can worsen an existing condition. In contrast, an adjustment disorder is a reaction to the stressor (e.g., having a medical illness).

Normative stress reactions. When bad things happen, most people get upset. This is not an adjustment disorder. The diagnosis should only be made when the magnitude of the distress (e.g., alterations in mood, anxiety, or conduct) exceeds what would normally be expected (which may vary in different cultures) or when the adverse event precipitates functional impairment.

Comorbidity

Adjustment disorders can accompany most mental disorders and any medical disorder. Adjustment disorders can be diagnosed in addition to another mental disorder only if the latter does not explain the particular symptoms that occur in reaction to the stressor. For example, an individual may develop an adjustment disorder, with depressed mood, after losing a job and at the same time have a diagnosis of obsessive-compulsive disorder. Or, an individual may have a depressive or bipolar disorder and an adjustment disorder as long as the criteria for both are met. Adjustment disorders are common accompaniments of medical illness and may be the major psychological response to a medical disorder.

Other Specified Trauma- and Stressor-Related Disorder

309.89 (F43.8)

This category applies to presentations in which symptoms characteristic of a trauma- and stressor-related disorder that cause clinically significant distress or impairment in social, occupational, or other important areas of functioning predominate but do not meet the full criteria for any of the disorders in the trauma- and stressor-related disorders diagnostic class. The other specified trauma- and stressor-related disorder category is used in situations in which the clinician chooses to communicate the specific reason that the presentation does not meet the criteria for any specific trauma- and stressor-related disorder. This is done by recording "other specified trauma- and stressor-related disorder" followed by the specific reason (e.g., "persistent complex bereavement disorder").

Examples of presentations that can be specified using the "other specified" designation include the following:

1. **Adjustment-like disorders with delayed onset of symptoms that occur more than 3 months after the stressor.**
 2. **Adjustment-like disorders with prolonged duration of more than 6 months without prolonged duration of stressor.**
 3. **Ataque de nervios:** See "Glossary of Cultural Concepts of Distress" in the Appendix.
 4. **Other cultural syndromes:** See "Glossary of Cultural Concepts of Distress" in the Appendix.
 5. **Persistent complex bereavement disorder:** This disorder is characterized by severe and persistent grief and mourning reactions (see the chapter "Conditions for Further Study").
-

Unspecified Trauma- and Stressor-Related Disorder

309.9 (F43.9)

This category applies to presentations in which symptoms characteristic of a trauma- and stressor-related disorder that cause clinically significant distress or impairment in social, occupational, or other important areas of functioning predominate but do not meet the full criteria for any of the disorders in the trauma- and stressor-related disorders diagnostic class. The unspecified trauma- or stressor-related disorder category is used in situations in which the clinician chooses *not* to specify the reason that the criteria are not met for a specific trauma- and stressor-related disorder, and includes presentations in which there is insufficient information to make a more specific diagnosis (e.g., in emergency room settings).
