

Introduction to Plasma Physics (PY5012)

Lectures 1 & 2: Basic Concepts






Dr. Peter T. Gallagher
Astrophysics Research Group
Trinity College Dublin


What is a plasma?

- “A plasma is a quasi-neutral gas consisting of positively and negatively charged particles (usually ions and electrons) which are subject to electric, magnetic and other forces, and which exhibit collective behaviour.”
 - Schwartz, Owen and Burgess, “Astrophysical Plasmas”.
- Langmuir in 1920s showed that an ionised gas can support oscillations, which resemble a jelly-like substance. He named it a “plasma”.
- Scientific term for “plasma” was introduced in 1839 by Czech biologist to describe a jelly-like medium of cells.

Plasmas in Nature and Technology



Hierarchy of plasma phenomena



Basic Parameters: Speed, Energy and Temperature

- For ensemble of N particles of mass m and velocity u , average energy per particle is

$$\langle E \rangle = \frac{1}{2N} \sum_{i=1}^N m_i u_i^2$$

- In thermal equilibrium, particles have Maxwell-Boltzmann distribution of speeds.

$$f(u) = N \sqrt{\frac{m}{2\pi k_B T}} e^{-1/2 mu^2 / k_B T}$$

- Average KE can be calculated using

$$\langle E \rangle = \frac{\int_{-\infty}^{+\infty} 1/2 mu^2 f(u) du}{\int_{-\infty}^{+\infty} f(u) du}$$

- Integrating numerator by parts, and denominator using $\int_{-\infty}^{+\infty} e^{-a^2 x^2} dx = \sqrt{\pi} / a$ we have $\langle E \rangle = 1/2 k_B T$ or in 3D,

$$\langle E \rangle = 3/2 k_B T$$

Basic Parameters: Speed, Energy and Temperature


- Using $\langle E \rangle = 3/2 k_B T$, average KE of a gas at 1 K is $\sim 2.07 \times 10^{-23}$ J.
- Plasma temperature often given in electron Volts (eV), where 1 eV = 1.602×10^{-19} Coulombs (C).
- For electron or proton, $|q| = 1.602 \times 10^{-19}$ C, therefore

$$E = \frac{1/2 mu^2}{2|q|} \text{ eV}$$

- **Note:** Typically, energy corresponding to $k_B T$ used for temperature, where $k_B T = 1 \text{ eV} = 1.602 \times 10^{-19}$ J.
- Q. Show that a 0.5 eV plasma corresponds to a temperature of 5,800 K (solar photosphere).

Range of plasma parameters


- Density range: >30 orders of magnitude. Temperature range: ~10 orders.



Plasma Oscillations and Plasma Frequency

- Consider plasma of equal number of positive and negative charges. Overall, plasma is neutral, so $n_e = n_i = n$

- Now displace group of electrons by Δx .
- Charge separation gives \mathbf{E} , which accelerates electrons towards initial position. Electrons overshoot equilibrium position.



- Using Newton's Law,
$$m_e \frac{d^2 \Delta x}{dt^2} = eE \quad (1)$$

- Displacement sets up E across distance L , similar to parallel plate capacitor.

- Charge per unit area of slab is $\sigma = -ne\Delta x \Rightarrow E = \sigma / \epsilon_0 = -ne\Delta x / \epsilon_0$

Plasma Oscillations and Plasma Frequency

- Therefore, Eqn. (1) can be written $\frac{d^2\Delta x}{dt^2} = -\frac{ne^2}{m_e\epsilon_0}\Delta x = -\omega_p^2\Delta x$

where

$$\omega_p^2 = \frac{ne^2}{m_e\epsilon_0} \quad \text{Electron Plasma Frequency}$$

- Plasma oscillations are result of plasma trying to maintain charge neutrality.
- Plasma frequency commonly written $f_p = \frac{\omega_p}{2\pi} = 9000\sqrt{n_e}$ Hz where n_e is in cm^{-3} .
- In Solar System, f_p ranges from hundreds of MHz (in solar corona) to <1 kHz (near outer planets).
- What is plasma frequency of Earth's ionosphere? What implication does this have for (i) short wave radio communications and (ii) radio astronomy?

Plasma Criteria

- In a partially ionized gas where collisions are important, plasma oscillations can only develop if mean free time between collisions (τ_c) is long compared with the oscillation period ($\tau_p = 1/\omega_p$).
- That is, $\tau_c \gg \tau_p$ or $\tau_c/\tau_p \gg 1$ *Plasma Criterion #1*
- Above is a *criterion* for an ionized gas to be considered a plasma.
- Behaves like neutral gas is criterion is not true.
- Plasma oscillations can be driven by natural thermal motions of electrons ($E=1/2k_B T_e$). Work by displacement of electron by Δx is and using $E = -ne\Delta x / \epsilon_0$

$$\begin{aligned} W &= \int F dx \\ &= \int_0^{\Delta x} eE(x) dx \\ &= \frac{e^2 n \Delta x^2}{2\epsilon_0} \end{aligned}$$

Plasma Criteria

- Equating work done by displacement with average energy in thermal agitation

$$\frac{e^2 n \Delta x^2}{2 \epsilon_0} \approx \frac{1}{2} k_B T_e$$

- The maximum distance an electron can travel is $\Delta x_{max} = \lambda_D$, where

$$\lambda_D = \sqrt{\frac{\epsilon_0 k_B T_e}{e^2 n}} \quad \text{Debye Length}$$

- Gas is considered a plasma if length scale of system is larger than Debye Length:

$$\lambda_D \ll L \quad \text{Plasma Criterion \#2}$$

- Debye Length is spatial scale over which charge neutrality is violated by spontaneous fluctuations.
- Debye Number is defined as $N_D = n [4\pi\lambda_D^3/3]$

Debye Shielding

- Plasmas do not contain strong electric fields as they reorganize to shield from them.
- Plasma oscillations are excited to assert *macroscopic* neutrality.
- If plasma subjected to external E , free charges redistribute so that plasma is shielded.
- Suppose immerse test particle $+Q$ within a plasma with $n_i = n_e = n$
- At $t = 0$, electric potential is $\Phi(r) = \frac{1}{4\pi\epsilon_0} \frac{Q}{r}$
- As time progresses, electrons are attracted, while ions are repelled. As $m_i \gg m_e$ neglect motion of ions.
- At $t \gg 0$, $n_e > n$ and a new potential is set up, with charge density:

$$\rho = e(n_e - n_i).$$

Debye Shielding

- New potential evaluated using Poisson's equation:

$$\nabla^2 \Phi(r) = -\frac{\rho}{\epsilon_0} = -\frac{e(n_e - n_i)}{\epsilon_0}$$

- In presence of potential, electron number density is

$$n_e(r) = n e^{-e\Phi(r)/k_b T_e}$$

- Subbing this into Poisson's equation in spherical coordinates.

$$\frac{1}{r^2} \frac{d}{dr} \left(r^2 \frac{d\Phi}{dr} \right) = -\frac{en}{\epsilon_0} \left[e^{-e\Phi/k_b T_e} - 1 \right]$$

- For $|e\Phi| \ll k_b T_e$ can us Taylor expansion: $e^x \approx 1 + x \Rightarrow$

$$\frac{1}{r^2} \frac{d}{dr} \left(r^2 \frac{d\Phi}{dr} \right) \approx \left[\frac{ne^2}{\epsilon_0 k_b T_e} \right] \Phi(r) = \frac{1}{\lambda_D^2} \Phi(r)$$

where λ_D is the *Debye shielding length*.

Debye Shielding

- Solution to previous is $\Phi(r) = \left[\frac{1}{4\pi\epsilon_0} \frac{Q}{r} \right] e^{-r/\lambda_D}$

- As $r \rightarrow 0$, potential is that of a free charge in free space, but for $r \gg \lambda_D$ potential falls exponentially.

- Coloumb force is long range in free space, but only extends to Debye length in plasma.

- For positive test charge, shielding cloud contains excess of electrons.

- Recall $\lambda_D = \sqrt{\frac{\epsilon_0 k_b T_e}{e^2 n}}$


\Rightarrow size of shielding cloud increases as electron temperature as electrons can overcome Coulomb attraction. Also, λ_D is smaller for denser plasma because more electrons available to populate shielding cloud.

Debye Shielding

- Useful numerical expression for Debye length:

$$\lambda_D \cong 69 \sqrt{T_e/n}$$

where T_e is in K and n is in m^{-3} .



The Plasma Parameter

- The typical number of particles in a Debye sphere is given by *the plasma parameter*:

$$\begin{aligned} \Lambda &= 4\pi n \lambda_D^3 \\ &= \frac{1.38 \times 10^6 T_e^{3/2}}{n^{1/2}} \end{aligned}$$

- Defined as N_D in Inan & Gollowski and usually given as argument of *Coulomb Logarithm* ($\log(\Lambda)$).
- If $\Lambda \ll 1$, the Debye sphere is sparsely populated, corresponding to a strongly coupled plasma.
- Strongly coupled plasmas tend to be cold and dense, whereas weakly coupled tend to be diffuse and hot.

The Plasma Parameter

Description	Plasma parameter magnitude	
	$\Lambda \ll 1$	$\Lambda \gg 1$
Coupling	Strongly coupled plasma	Weakly coupled plasma
Debye sphere	Sparsely populated	Densely populated
Electrostatic influence	Almost continuously	Occasional
Typical characteristic	Cold and dense	Hot and diffuse
Examples	Solid-density laser ablation plasmas Very "cold" "high pressure" arc discharge Inertial fusion experiments White dwarfs / neutron stars atmospheres	Ionospheric physics Magnetic fusion devices Space plasma physics Plasma ball

Collisions

- Neutral particles have quite small collision cross sections. As Coulomb force is long range, charged particles are more frequent.

- A *collisional plasma* is one where $\lambda_{mfp} \ll L$

where L is the observational length scale and $\lambda_{mfp} = 1/\sigma n$ is the mean free path.

- The effective Coulomb cross-section is $\sigma = \pi r_c^2$
- An electron will be affected by a neighbouring ion if the Coulomb potential is of the order of the electron thermal energy:

$$\frac{e^2}{4\pi\epsilon_0 r_c} \approx \frac{3}{2} k_b T$$

$$\Rightarrow \sigma = \pi \left(\frac{e^2}{6\pi k_b \epsilon_0} \right)^2 \frac{1}{T^2}$$

- At $T = 10^6$ K, $\sigma \sim 10^{-22}$ m², which is much larger than the geometric nuclear cross-section of 10^{-33} m².

Collisions

- In terms of the plasma parameter, the collision frequency ($\nu = n e \nu$) is

$$\nu \approx \frac{\omega_p}{64\pi} \frac{\ln \Lambda}{\Lambda}$$

where $\ln(\Lambda)$ is the Coulomb logarithm. Used as Λ is large, but $10 < \ln(\Lambda) < 30$.

- In a weakly coupled plasma, $\nu \ll \omega_p \Rightarrow$ collisions do not affect plasma oscillations
- More rigorously, it can be shown that

$$\nu \approx \frac{\sqrt{2}\omega_p^4}{64\pi n_e} \left(\frac{k_B T}{m_e} \right)^{-3/2} \ln \Lambda$$

- Thus, diffuse, high temperature plasmas tend to be collisionless. See page 156 of Inan & Golkowski.

Examples of key plasma parameters

	$n(\text{m}^{-3})$	$T(\text{eV})$	$\omega_p(\text{sec}^{-1})$	$\lambda_D(\text{m})$	Λ
Interstellar	10^6	10^{-2}	6×10^4	0.7	4×10^6
Solar Chromosphere	10^{18}	2	6×10^{10}	5×10^{-6}	2×10^3
Solar Wind (1AU)	10^7	10	2×10^5	7	5×10^{10}
Ionosphere	10^{12}	0.1	6×10^7	2×10^{-3}	1×10^5
Arc discharge	10^{20}	1	6×10^{11}	7×10^{-7}	5×10^2
Tokamak	10^{20}	10^4	6×10^{11}	7×10^{-5}	4×10^8
Inertial Confinement	10^{28}	10^4	6×10^{15}	7×10^{-9}	5×10^4

Table 1.1: Key parameters for some typical weakly coupled plasmas.

* From *Plasma Physics* by R. Fitzpatrick

Next Lecture: Single-particle Motion

