

FLESH FLY

Scientific name: *Sarcophaga haemorrhoidalis* (Fallén)

Family: Sarcophagidae

Order: Diptera

Introduction

Sarcophaga haemorrhoidalis (Fallén), commonly known as the red-tailed flesh fly, is a member of the family Sarcophagidae, the flesh flies. Flesh flies are calypteres possessing well-developed lower calypteres which are associated with the wings. Red-tailed flesh flies are considered opportunistic feeders on carrion, and primarily feed on human and animal waste (Madubunyi 1986). The flesh flies are an important family of insects in forensic entomology, a discipline that focuses on arthropods and arthropod behavior as evidence in legal and criminal matters.



Figure 1. Dorsal view of adult *Sarcophaga haemorrhoidalis* (Fallén), the red-tailed flesh fly. Photograph by Jerry Butler, University of Florida.

Description

Adults: The adults of *S. haemorrhoidalis* resemble house flies (Family Muscidae) in their general appearance, but are larger and more robust, ranging from 8 to 14 mm in length. Adult *S. haemorrhoidalis* possess black-colored bodies covered with a whitish powder-like substance that gives them a grayish tint. Their abdomen has a distinct checkerboard-like pattern. The common name of this species comes from the presence of red-colored external genitalia visible at the terminal end of the abdomen. Like many other sarcophagids, they have large compound eyes and the arista of the antennae are long and are plumose at the base (Byrd and Castner 2001). The

adult head has a row of frontal bristles near the enlarged median vertical area. The mouthparts are designed for lapping liquid media from sugar and decomposing sources.



Figure 2. Lateral view of an adult *Sarcophaga haemorrhoidalis* (Fall n), the red-tailed flesh fly. Photograph by [Donald Hall](#), University of Florida.

Eggs: The eggs of *S. haemorrhoidalis* range in size from 0.5 to 3.5 mm long and 0.12 to 0.8 mm wide.

Larvae: The larvae are white to slightly yellow in color, and worm-like with a pointed head. The larvae are comparatively larger than other fly larvae, reaching 10 to 22 mm in length prior to pupation (Baker et al. 1986). Each instar has a pair of strong mandibles for tearing and eating flesh and other food substrates (Mullen and Durden 2002).

Pupae: The pupae are reddish-orange to dark brown and change colors as they age.

Life Cycle

S. haemorrhoidalis undergoes complete metamorphosis through four life stages: egg, larva, pupa, and adult. The complete life cycle takes approximately 23 days under laboratory conditions from 1st instar larvae to adult (Madubunyi 1986). Like other sarcophagids, females engage in larviposition, wherein the eggs hatch inside the female and are then deposited as first instar maggots on decomposing flesh or excrement (Byrd and Castner 2001).



Figure 3. Adult female *Sarcophaga haemorrhoidalis* (Fall n), the red-tailed flesh fly, depositing first instar maggots (white larvae, right). Photograph by [Jerry Butler](#), University of Florida.

Larvae prefer moist environments and can even survive in semi-aquatic conditions (Byrd 1998-2011). On average, larvae complete three instars within five to six days before pupating (Smith and Whitman 1992), though other studies have shown larval duration to be seven to nine days at 25C° (Pape 1996). The reason for these differences is that the larvae are greatly affected by temperature, with higher temperatures generally accelerating larval development, especially during the third instar.

Pupae develop more quickly at cool temperatures and are on average 12 mm in length (Byrd and Butler 1998) and the entire pupal stage can take approximately 16 days (Smith and Whitman 1992). Within two to nine days after adults emerge from the puparia, copulation occurs. There is no known courtship ritual for this species; males have no specialized ability to identify females of the species. In fact, it has been noted that males will copulate with other males and dead flies. Additionally, the same male has been observed mating with the same female multiple times suggesting that this species does not have a pheromone or other cues to detect previous insemination as has been observed for other Dipteran species.



Figure 4. *Sarcophaga haemorrhoidalis* (FallÅ©n), red-tailed flesh flies, mating. Photograph by [Donald Hall](#), University of Florida.

After insemination, eggs are retained in the female and require a minimum incubation of nine days with an average of 13 days before larviposition occurs (Madubunyi 1986). Females can lay 40 to 80 first instar larvae per reproductive cycle (Braverman et al. 1994).

Medical and Economic Importance

Sarcophaga haemorrhoidalis is of medical importance as it is one of 20 species known to cause gastrointestinal myiasis and the infestation of living tissue (Mullen and Durden 2002). Myiasis can occur in livestock, including ungulates, which can result in economic loss. The main reason that *S. haemorrhoidalis* is a myiasis-causing species is due to its attraction to feces and preference to larviposit indoors (Byrd and Castner 2001). Myiasis most often occurs from ingesting contaminated food and infestations may be persistent if left untreated (Smith &

Whitman 1992). Myiasis is most likely to occur in areas with overcrowding and hygienic conditions that are substandard (Braverman et al. 1994) but can also occur in hospitals and nursing homes.

Adult *S. haemorrhoidalis* have been connected with the transmission of other disease organisms including *Escherichia coli* (Migula), *Shigella dysenteriae* (Shiga), *Streptococcus* spp., *Salmonella* spp., tapeworms, and even the virus responsible for polio (Smith and Whitman 1992).

Forensic Importance

Sarcophaga haemorrhoidalis is particularly important to forensic entomology as it can be the first, or one of the first, arthropods to arrive at a corpse. The species is a strong flyer and has the ability to fly in inclement weather when other arthropod species are unable to fly. Like blow fly species, sarcophagid flies are also used to determine the post-mortem interval (PMI) in death investigations (Byrd and Butler 1998), and because their development cycle is short, they are the most useful for the first three to four weeks after death.

Management

Control of *S. haemorrhoidalis* is the same as for blow flies, house flies and other filth flies. Elimination of suitable food substrates by bagging trash and other decaying material and covering garbage containers will limit infestations by this species. Using mesh screens at doorways and windows as well as using air currents at speeds of 1500 ft/min (450 m/min) will also reduce access to indoor areas. Pesticides can be used to eliminate indoor *S. haemorrhoidalis* infestations using space sprays and residual sprays that are designed to control adult flies (Ebeling 2002).