

SOIL MICROBIOLOGY

KEY POINTS

- Biological fertility is under-studied and our scientific knowledge of it is incomplete.
- In addition to fertility, soil microorganisms also play essential roles in the nutrient cycles that are fundamentally important to life on the planet.
- In the past, agricultural practices have failed to promote healthy populations of microorganisms, limiting production yields and threatening sustainability.
- Scientific research is exploring new and exciting possibilities for the restoration and promotion of healthy microbial populations in the soil.

'Soil is essential for the maintenance of biodiversity above and below ground. The wealth of biodiversity below ground is vast and unappreciated: millions of microorganisms live and reproduce in a few grams of topsoil, an ecosystem essential for life on earth...'

Soil has traditionally been seen as a "dead" agricultural medium – something to keep crops upright. Soil physicochemical analyses were conducted to determine the application rate of chemical fertilizers to sustain/increase the season's yields. Soils were injudiciously ploughed, drenched in herbicides and pesticides, crop residues were burned, and fields were left bare and vulnerable to nature's elements, with precious fertile topsoil being blown away or carried off during wind and rainstorms. The vital role of soil microorganisms in agriculture have only recently gained popularity with South African farmers after they experimented with various agricultural practices in an effort to increase their yields and soil's health/fertility in a sustainable way. Before this paradigm shift, we never realized that only 6-8 cm of fertile soil was naturally formed over a period of 2000 years; that soil-life consisted of thousands of different insects, earthworms, mites, nematodes, fungi, yeasts, and single-cell organisms; that a teaspoon of fertile soil could contain a billion bacteria and almost 5,000 different species of bacteria per gram of soil.

After decades of collaboration between researchers and farmers, conservation agriculture (CA) was promoted as the most probable solution to sustainable agriculture. Conservation agriculture aims to improve/sustain productivity, increase profits and food security while preserving and enhancing the resource base and the environment. Long-term implementation of CA's three main principles, i.e. minimum soil disturbance, permanent organic soil cover, and crop diversification, will inevitably lead to healthier soil and sustainable agriculture.

Nematodes, fungi and bacteria are usually associated with large-scale yield-losses, but what we fail to realize in a healthy soil with its high microbial diversity and activity, is that beneficial and harmful organisms are present in a very fine balance. With this sensitive balance maintained, plant-pathogens are suppressed and/or out-competed by the indigenous soil life, and vital soil processes are optimally executed. This attribute is highly beneficial, especially when external forces, such as drought and diseases, disrupt soil processes performed by specific species. In the event of such an external force, these vital processes can immediately be re-initiated and

maintained by other individual or groups of species, thus strengthening the soil's resilience and resistance to disruptive forces.

Due to the sensitive nature of soil microbial populations, they could be used as "early warning systems" or sentinel organisms to detect deterioration or improvement in soil quality – almost like the canaries that were historically carried into the coal mine tunnels by the miners to detect the collection of dangerous gases. If the canary was killed by the gases, it served as a warning to coal miners to exit the tunnels immediately. With the case of soil microbial populations, for example, when a soil's microbial diversity (the number of different microorganisms) decreases but the activity (how hard/fast they work) increases, or *vice versa*, it indicates a disturbance in the soil's balance and that the soil's health might be compromised. Such an imbalance typically occurs during injudicious ploughing and fertiliser application, extended fallow periods, or continuous monocropping.

Soil microbiology is the study of organisms in soil, their functions and how they affect soil properties. Soil microorganisms can be classified as bacteria, actinomycetes, fungi, algae, protozoa and viruses. Each of these groups has different characteristics that define the organisms and different functions in the soil it lives in. Importantly, these organisms do not exist in isolation; they interact and these interactions influence soil fertility as much or more than the organism's individual activities.




Figure 1: Colonies of soil bacteria (shown in light blue). Each bacterium is approximately one micron in size.

Source: Karl Ritz (soilquality.org.au)

Bacteria: Bacteria are organisms that have only one cell and are, therefore, microscopic. There are anywhere from 100 million to one billion bacteria in just a teaspoon of moist, fertile soil. They are decomposers, eating dead plant material and organic waste. By doing this, the bacteria release nutrients that other organisms could not access. The bacteria do this by changing the nutrients from inaccessible to usable forms. The process is essential in the nitrogen cycle.

Actinomycetes: Actinomycetes are soil microorganisms like both bacteria and fungi, and have characteristics linking them to both groups. They are often believed to be the missing evolutionary

link between bacteria and fungi, but they have many more characteristics in common with bacteria than they do fungi. Actinomycetes give soil its characteristic smell. They have also been the source of several significant therapeutic medicines.




Figure 2: Fungal filaments, or hyphae, (shown in blue) extending through soil.
Source: Karl Ritz (soilquality.org.au)

Fungi: Fungi are unusual organisms, in that they are not plants or animals. They group themselves into fibrous strings called hyphae. The hyphae then form groups called mycelium which are less than 0.8mm wide but can get as long as several metres. They are helpful, but could also be harmful, to soil organisms. Fungi are helpful because they have the ability to break down nutrients that other organisms cannot. They then release them into the soil, and other organisms get to use them. Fungi can attach themselves to plant roots. Most plants grow much better when this happens. This is a beneficial relationship called mycorrhizal. The fungi help the plant by giving it needed nutrients and the fungi get carbohydrates from the plant, the same food that plants give to humans. On the other hand, fungi can get food by being parasites and attaching themselves to plants or other organisms for selfish reasons.

Some of the functions performed in soil by fungi are:

- **Decomposers** – saprophytic fungi – convert dead organic material into fungal biomass, carbon dioxide (CO₂), and small molecules, such as organic acids.
- **Mutualists** – the mycorrhizal fungi – colonise plant roots. In exchange for carbon from the plant, mycorrhizal fungi help to make phosphorus soluble and bring soil nutrients (phosphorus, nitrogen, micronutrients and, perhaps, water) to the plant. One major group of mycorrhizae, the *ectomycorrhizae*, grow on the surface layers of the roots and are commonly associated with trees. The second major group of mycorrhizae are the *endomycorrhizae* that grow within the root cells and which are commonly associated with grasses, row crops, vegetables and shrubs.
- **Parasites:** The third group of fungi, *pathogens* or *parasites*, causes reduced production or death when they colonise roots and other organisms.

Algae: Algae are present in most of the soils where moisture and sunlight are available. Their number in the soil usually ranges from 100 to 10,000 per gram of soil. They are capable of photosynthesis, whereby they obtain carbon dioxide from atmosphere and energy from sunlight and synthesise their own food.

The major roles and functions of algae in soil are:

- Playing an important role in the maintenance of soil fertility, especially in tropical soils.
- Adding organic matter to soil when they die and thus increasing the amount of organic carbon in soil.
- Acting as a cementing agent by binding soil particles and thereby reducing and preventing soil erosion.
- Helping to increase the water retention capacity of soil for longer time periods.
- Liberating large quantities of oxygen in the soil environment through the process of photosynthesis and, thus, facilitating submerged aeration.
- Helping to check the loss of nitrates through leaching and drainage, especially in un-cropped soils.
- Helping in the weathering of rocks and the building up of soil structure.

Protozoa: These are colourless, single-celled animal-like organisms. They are larger than bacteria, varying from a few microns to a few millimetres. Their population in arable soil ranges from 10,000 to 100,000 per gram of soil and they are abundant in surface soil. They can withstand adverse soil conditions, as they are characterised by a protected, dormant stage in their life cycle.

The major functions, roles and features of protozoa are:

- Most protozoans derive their nutrition from feeding or ingesting soil bacteria and, thus, they play an important role in maintaining microbial/bacterial equilibrium in the soil.
- Some protozoa have been recently used as biological control agents against organisms that cause harmful diseases in plants.
- Several soil protozoa cause diseases in human beings that are carried through water and other vectors. Amoebic dysentery is an example.

Viruses: Soil viruses are of great importance, as they may influence the ecology of soil biological communities through both an ability to transfer genes from host to host and as a potential cause of microbial mortality. Consequently, viruses are major players in global cycles, influencing the turnover and concentration of nutrients and gases.

Despite this importance, the subject of soil virology is understudied. To explore the role of the viruses in plant health and soil quality, studies are being conducted into virus diversity and abundance in different geographic areas (ecosystems). It has been found that viruses are highly abundant in all the areas studied so far, even in circumstances where bacterial populations differ significantly in the same environments. Soils probably harbour many novel viral species that, together, may represent a large reservoir of genetic diversity. Some researchers believe that investigating this largely unexplored diversity of soil viruses has the potential to transform our understanding of the role of viruses in global ecosystem processes and the evolution of microbial life itself.



Figure 3: A soil nematode worm. Harmful species have received much attention, but far less is known about the majority of the nematode community, which play beneficial roles in soil.
Source: Karl Ritz (soilquality.org.au)

Nematodes: Not microorganisms (strictly speaking), nematode worms are typically 50 microns in diameter and one millimetre in length. Species responsible for plant diseases have received much attention, but far less is known about much of the nematode community, which play beneficial roles in soil. An incredible variety of nematodes have been found to function at several levels of the soil food web. Some feed on the plants and algae (the first level), others are grazers that feed on bacteria and fungi (second level), and some feed on other nematodes (higher levels).

Free-living nematodes can be divided into four broad groups based on their diet. Bacterial-feeders consume bacteria. Fungal-feeders feed by puncturing the cell walls of fungi and sucking out the internal contents. Predatory nematodes eat all types of nematodes and protozoa. They eat smaller organisms whole or attach themselves to the cuticle of larger nematodes, scraping away until the prey's internal body parts can be extracted. Like protozoa, nematodes are important in mineralising, or releasing, nutrients in plant-available forms. When nematodes eat bacteria or fungi, ammonium is released because bacteria and fungi contain much more nitrogen than the nematodes require.

Nematodes may also be useful indicators of soil quality because of their tremendous diversity and their participation in many functions at different levels of the soil food web.

Role and Functions

Collectively, soil microorganisms play an essential role in decomposing organic matter, cycling nutrients and fertilising the soil. Without the cycling of elements, the continuation of life on Earth would be impossible, since essential nutrients would rapidly be taken up by organisms and locked in a form that cannot be used by others. The reactions involved in elemental cycling are often chemical in nature, but biochemical reactions, those facilitated by organisms, also play an important part in the cycling of elements. Soil microbes are of prime importance in this process. Soil microbes are also important for the development of healthy soil structure. Soil microbes produce lots of gummy substances (polysaccharides and mucilage, for example) that help to cement soil aggregates. This cement makes aggregates less likely to crumble when exposed to

water. Fungal filaments also stabilise soil structure because these threadlike structures branch out throughout the soil, literally surrounding particles and aggregates like a hairnet. The fungi can be thought of as the “threads” of the soil fabric. It must be stressed that microbes generally exert little influence on changing the actual physical structure of the soil; that is performed by larger organisms.

Soil microorganisms are both components and producers of soil organic carbon, a substance that locks carbon into the soil for long periods. Abundant soil organic carbon improves soil fertility and water-retaining capacity. There is a growing body of research that supports the hypothesis that soil microorganisms, and fungi in particular, can be harnessed to draw carbon out of the atmosphere and sequester it in the soil. Soil microorganisms may provide a significant means of reducing atmospheric greenhouse gasses and help to limit the impact of greenhouse gas-induced climate change.