



Food and Agriculture
Organization of the
United Nations



Government of Pakistan
Planning Commission
Nutrition Section

PAKISTAN DIETARY GUIDELINES FOR BETTER NUTRITION

My Plate



Published by
Food and Agriculture Organization of the United Nations
and
Ministry of Planning Development and Reform, Government of Pakistan

Pakistan Dietary Guidelines for Better Nutrition

**Published by
Food and Agriculture Organization of the United Nations
and
Ministry of Planning Development and Reform, Government of Pakistan**

The designations employed and the presentation of material in this information product do not imply the expression of any opinion whatsoever on the part of the Food and Agriculture Organization of the United Nations (FAO) or the Ministry of Planning Development and Reform, Government of Pakistan, concerning the legal or development status of any country, territory, city or area or of its authorities, or concerning the delimitation of its frontiers or boundaries. The mention of specific companies or products of manufacturers, whether or not these have been patented, does not imply that these have been endorsed or recommended by FAO or the Ministry in preference to others of a similar nature that are not mentioned.

The views expressed in this information product are those of the author(s) and do not necessarily reflect the views or policies of FAO or the Ministry.

ISBN 978-92-5-131021-2 (FAO)

© FAO, 2018



Some rights reserved. This work is made available under the Creative Commons Attribution-NonCommercial-ShareAlike 3.0 IGO licence (CC BY-NC-SA 3.0 IGO; <https://creativecommons.org/licenses/by-nc-sa/3.0/igo>).

Under the terms of this licence, this work may be copied, redistributed and adapted for non-commercial purposes, provided that the work is appropriately cited. In any use of this work, there should be no suggestion that FAO endorses any specific organization, products or services. The use of the FAO logo is not permitted. If the work is adapted, then it must be licensed under the same or equivalent Creative Commons license. If a translation of this work is created, it must include the following disclaimer along with the required citation: "This translation was not created by the Food and Agriculture Organization of the United Nations (FAO). FAO is not responsible for the content or accuracy of this translation. The original English edition shall be the authoritative edition."

Any mediation relating to disputes arising under the licence shall be conducted in accordance with the Arbitration Rules of the United Nations Commission on International Trade Law (UNCITRAL) as at present in force.

Third-party materials. Users wishing to reuse material from this work that is attributed to a third party, such as tables, figures or images, are responsible for determining whether permission is needed for that reuse and for obtaining permission from the copyright holder. The risk of claims resulting from infringement of any third-party-owned component in the work rests solely with the user.

Sales, rights and licensing. FAO information products are available on the FAO website (www.fao.org/publications) and can be purchased through publications-sales@fao.org. Requests for commercial use should be submitted via: www.fao.org/contact-us/licence-request. Queries regarding rights and licensing should be submitted to: copyright@fao.org.

CONTENTS

| | |
|---|------|
| FOREWORD | vii |
| ACKNOWLEDGEMENTS | viii |
| MESSAGE FROM THE FAO REPRESENTATIVE | ix |
| ACRONYMS | x |
| EXECUTIVE SUMMARY | xi |
| INTRODUCTION | 1 |
| Rationale | 2 |
| Goals..... | 3 |
| Objectives..... | 3 |
| NUTRITION SITUATION | 4 |
| Nutritional status of infants (0-6 months)..... | 4 |
| Nutritional status of infants and young children (6-24 months)..... | 4 |
| Nutritional status of children (24-36 months)..... | 5 |
| Nutritional status of children (3-10 years) | 6 |
| Nutritional status of adolescents (10-19 years) | 7 |
| Nutritional status of adults (19-60 years) | 8 |
| Nutritional status of elderly (>60 years)..... | 8 |
| Nutritional status of pregnant women..... | 9 |
| Nutritional status of lactating women | 10 |
| Food availability and consumption in Pakistan | 11 |
| Relationship between diet and disease | 15 |
| Traditional and fast foods | 18 |
| Glycemic index of foods | 19 |
| Food allergy..... | 19 |
| Unprocessed and processed foods | 19 |
| Nutrition labelling | 20 |
| Overweight and obesity | 21 |
| Physical activity and lifestyle behaviour..... | 22 |
| Smoking and smokeless tobacco use | 23 |

| | |
|---|----|
| Food safety and hygiene practices | 24 |
| hygienic food management..... | 24 |
| Water, sanitation and hygiene | 24 |
| METHODOLOGY..... | 26 |
| PAKISTAN DIETARY GUIDELINES FOR BETTER NUTRITION (PDGN) | 26 |
| Food and food groups | 27 |
| Cereals..... | 27 |
| Vegetables | 28 |
| Fruits..... | 28 |
| Milk and milk products..... | 29 |
| Meat and pulses | 29 |
| Fats and oils..... | 29 |
| Recommended dietary allowances | 30 |
| Infants (0-6 months)..... | 30 |
| Infants and young children (6-24 months) | 31 |
| Children (24-36 months) | 31 |
| Children (3-10 years) | 32 |
| Adolescents (10-19 years) | 32 |
| Adults (19-60 years) | 32 |
| Elderly (>60 years)..... | 32 |
| Pregnancy..... | 33 |
| Lactation..... | 33 |
| MYPLATE | 34 |
| Portion sizes and frequency of servings..... | 35 |
| Dietary recommendations for infants and young children (6-24 months)..... | 36 |
| Food and portion sizes for children (24-36 months) | 37 |
| Food and portion sizes for children (3-10 years)..... | 38 |
| Food and portion sizes for adolescents (10-19 years)..... | 40 |
| Food and portion sizes for adults (19-60 years) | 41 |
| Food and portion sizes for elderly (>60 years) | 42 |

| | |
|---|----|
| Food and portion sizes for pregnant women | 44 |
| Food and portion sizes for lactating women | 45 |
| SAMPLE MENUS BY AGE AND PHYSIOLOGICAL STATUS..... | 46 |
| Sample menu for children (3-10 years) | 46 |
| Sample menu for adolescents (10-19 years)..... | 47 |
| Sample menu for adults (19-60 years) | 48 |
| Sample menu for elderly (> 60)..... | 49 |
| Sample menu for pregnant women | 50 |
| Sample menu for lactating women | 51 |
| Cost of healthy diets..... | 52 |
| DIETARY MESSAGES..... | 53 |
| Maintain normal body weight by consuming all food groups and performing regular physical activity | 53 |
| Half of your daily cereals intake should include whole grains..... | 53 |
| Eat five servings of fresh vegetables and fruits a day..... | 53 |
| Take two to three servings of milk and milk products in a day | 53 |
| Consume meat and meat products, fish and eggs in moderation..... | 54 |
| Encourage consumption of pulses to attain healthy growth | 54 |
| Consume fortified flour, grains and their products..... | 54 |
| Limit consumption of edible oil and fat in cooking | 54 |
| Reduce sugar intake, and limit intake of soft drinks, confectionaries, bakery products and commercial fruit drinks | 54 |
| Limit salt in cooking and always use iodized salt..... | 55 |
| Limit consumption of fatty foods and highly processed foods..... | 55 |
| Change sedentary lifestyle to physically active lifestyle..... | 55 |
| Exclusively breastfeed the baby in the first six months and continue breastfeeding along with complementary feeding at least for two years..... | 55 |
| Women should increase intake of all the food groups daily, especially foods that are rich in iron and take extra care during pregnancy and lactation..... | 56 |
| Drink plenty of water each day | 56 |
| Read nutrition labeling on packaged food products | 56 |

| | |
|--|----|
| ANNEXURE-I: Glycemic index and glycemic load of foods | 57 |
| ANNEXURE-II: Personal hygiene and food safety guidelines..... | 58 |
| ANNEXURE-III: Foods rich in omega fatty acids..... | 59 |
| ANNEXURE-IV: Recommended daily allowances for Pakistani population for selected major nutrients..... | 60 |
| ANNEXURE-V: Recommendations for infant and young child feeding..... | 61 |
| ANNEXURE-VI: Complementary foods for infants > 6 months..... | 62 |
| ANNEXURE-VII: Food and portion sizes for different age groups..... | 65 |
| ANNEXURE-VIII: Daily recommended intakes for energy and nutrients | 66 |
| ANNEXURE-IX: Food as a source of nutrients..... | 67 |
| ANNEXURE-X: Nutrients rich foods | 68 |
| ANNEXURE-XI: Healthy snacks | 71 |
| ANNEXURE-XII: Recommendations on physical activity for different age groups..... | 72 |
| ANNEXURE-XIII. Committees for Pakistan Dietary Guidelines for Better Nutrition | 73 |
| GLOSSARY | 74 |
| REFERENCES | 76 |

FIGURES

| | |
|---|----|
| Figure 1. Availability of cereals over different time periods | 12 |
| Figure 2: Availability of fats and cooking oils over different time periods..... | 12 |
| Figure 3. Availability of maize and sugar over different time periods..... | 13 |
| Figure 4. Availability of pulses, fruits and vegetables over different time periods | 13 |
| Figure 5. Per capita GDP from 1960-2015 | 14 |
| Figure 6. Poverty based on low calories consumption over the years..... | 14 |
| Figure 7. Calories contribution of different foods in poor and non-poor diets..... | 16 |
| Figure 8. Food consumption patterns of households..... | 16 |
| Figure 9. Frequency of food consumption based on 7 days dietary recall..... | 17 |
| Figure 10. Trends in food prices over the years | 17 |

FOREWORD

The Pakistan Dietary Guidelines for Better Nutrition (PDGN) are comprehensive country specific guidelines for the general public to adopt healthy eating practices, and prevent and reduce the risk of infectious and chronic diseases. These guidelines are simple to adopt, provide age and gender appropriate and culturally acceptable options to choose nutritious foods.

The PDGN provide a thorough review of the food security, health and nutrition situation of the population and is cognizant of the fact that about half of the population is deficient in one or more of essential nutrients, reflecting unhealthy dietary practices compounded by poor hygiene and sanitation. Women and young children are more vulnerable to nutritional deficiencies, morbidity and mortality due to their compromised health and nutritional status. The PDGN highlight the predominant consumption of energy dense foods and monotonous diets by majority of the population.

The emphasis of the PDGN is on the consumption of a variety of safe and nutritious foods including milk and milk products, meat and pulses, wholegrain cereals, vegetables and fruits and decreasing consumption of energy dense foods such as deep fried foods, bakery products, processed foods and reducing the amount of fat specifically saturated fat, oil, sugar and salt in cooking, as there is a strong relationship between diet and disease, specially non-communicable diseases such as diabetes, cardiovascular diseases and others that are associated with poor dietary habits. Moreover, physical activity is critical for all population groups to remain physically fit and healthy.

The PDGN are a rich resource for general public, decision and policy makers, planners, researchers, academicians, food and pharmaceutical industries, hospitals, agriculture extension and allied health professionals for integration into their own plan of actions and interventions. These guidelines will benefit communities, clients and consumers in making healthy food choices and well suited to break the vicious cycle of malnutrition when integrated into nutrition sensitive and the social development programmes of the country.

I appreciate the efforts of the contributors and hope that the use of these guidelines, would serve as an effective tool for improving nutrition and well being of the population.



Sartaj Aziz

Deputy Chairman

Planning Commission of Pakistan

28 May, 2018

ACKNOWLEDGEMENTS

The Ministry of Planning Development and Reform, gratefully acknowledges the generous support of the Food and Agriculture Organization of the United Nations and Scaling up Nutrition (SUN) Movement for developing the Pakistan Dietary Guidelines for Better Nutrition (PDGN). Mr. Francisco Gamarro (Late), Ex-Deputy Representative FAO, and Mr. Nasar Hayat, Assistant Representative FAO, deserve appreciation for their continuous support in accomplishing the task of developing PDGN.

The efforts of Dr. Parvez Iqbal Paracha, Nutrition Specialist in preparing the document and great contributions of Dr. Nomeena Anis, Nutritionist and Gender Focal Person, FAO, Islamabad in developing a comprehensive document are highly appreciated. The dedicated efforts, overall coordination and commitment of the Nutrition Section/ SUN Secretariat are commendable. Special thanks to Dr. M. Azeem Khan, Member Food Security and Climate Change, Planning Commission for his overall guidance in finalizing of this document. The Nutrition Section appreciates the valuable contribution of Dr. Mubrik Ali, Ex-Member Food Security and Climate Change, Planning Commission Pakistan.

The Ministry of Planning Development and Reform is highly indebted to distinguished and renowned academicians, researchers, programme managers, policy makers, development partners and other stakeholders for their valuable inputs in formulation of PDGN, which will serve as a source of guidance for planning and implementation of nutrition related programmes aimed at promoting healthy diets and preventing malnutrition and diseases.



M. Aslam Shaheen

Chief Nutrition/SUN Focal Person

Ministry of Planning Development & Reform Pakistan


MESSAGE FROM THE FAO REPRESENTATIVE

Pakistan Dietary Guidelines for Better Nutrition (PDGN) are developed for the general public to support healthy eating, improve overall health and reduce malnutrition in all its form, including preventing disease and premature deaths. It is an opportune time to disseminate the dietary guidelines, when the country is confronted with challenges of diet associated diseases including overweight, obesity, diabetes, hypertension, cardiovascular diseases and cancers.

PDGN are intended for use by public health and nutrition service providers, food and agriculture professionals, programmes managers and policy makers to frame cost-effective, high impact policies and interventions that would help the public in making better food choices that are safe, nutrient rich and healthy.

PDGN provide an affordable and sustainable means of improving nutrition and health wellbeing of the population, which is a pre-requisite for socio-economic development, peace, prosperity and security in the country. The cost of malnutrition and associated diseases outweigh the nominal cost of food therefore food and nutritional wellbeing should be prioritized in household expenditures.

Dissemination and use of dietary guidelines is essential to create and promote nutrition awareness among the population. Strong public and private partnerships and community involvement are critical for successful adoption of these dietary guidelines. I appreciate the efforts and contributions of the technical experts for the timely production of the PDGN, benefitting all segments of the society including the poor and rich as well as urban and rural with adoption of healthy eating practices to enhance the quality of life.



Minà Dowlatchahi
FAO Representative
Pakistan

ACRONYMS

| | |
|-------|---|
| AMDR | Acceptable macronutrient distribution range |
| BMI | Body mass index |
| BMR | Basal metabolic rate |
| CAD | Coronary artery disease |
| CVDs | Cardiovascular diseases |
| DALYs | Disability adjusted life years |
| DNA | Deoxyribonucleic acid |
| FAO | Food and Agriculture Organization of the United Nations |
| DGs | Dietary guidelines |
| FCPs | Food consumption patterns |
| HDL | High density lipoproteins |
| IYCF | Infant and young child feeding |
| Kcals | Kilocalories |
| LDL | Low density lipoproteins |
| MDER | Minimum dietary energy requirement |
| MPDR | Ministry of Planning, Development and Reform |
| NCDs | Non-communicable diseases |
| NNS | National Nutrition Survey |
| NoU | Number of undernourished |
| NRVs | Nutrient reference values |
| PBS | Pakistan Bureau of Statistics |
| PDGN | Pakistan dietary guidelines for better nutrition |
| PoU | Prevalence of undernourishment |
| RDAs | Recommended dietary allowances |
| SoFI | State of food insecurity |
| WASH | Water, sanitation and hygiene |
| WHO | World Health Organization |

EXECUTIVE SUMMARY

Pakistan Dietary Guidelines for Better Nutrition (PDGN) are dietary recommendations based on scientific knowledge related to nutritional requirements, food consumption patterns and nutrients intake of the population. PDGN are developed taking into account the country's food and nutrition situation and the role of diet in health and disease. Pakistan has been passing through a demographic, economic and nutrition transition characterized by rising population, food accessibility issues, unhealthy eating trends, lack of safe food handling and management practices, and sedentary lifestyles. This has led to an increase in nutritional disorders and diet related diseases such as diabetes, hypertension, cardiovascular diseases, cancers and infectious diseases in children, adolescents and adults, especially women.

First country specific Dietary Guidelines (DGs) were developed in 2005. Since then, new scientific evidence related to the beneficial effects of food and bioactive factors, harmful effects of adulterants, contaminants, pesticides and insecticides, unhealthy food consumption patterns and excessive use of salt, sugars, saturated and trans-fatty acids on health, has been generated. These scientific developments have been considered during the revision of PDGN. Consultative workshops were held, with the active participation of technical experts and relevant stakeholders to develop and finalize the framework of PDGN. An extensive literature search and desk review was undertaken to review the food consumption patterns, nutrients intake and nutritional and health status of the population. National and international recommended dietary allowances (RDAs) and dietary guidelines were reviewed to extract relevant information for the development of PDGN.

Diet is one of the strongest determinants of health and nutritional status. Inappropriate and unhealthy dietary practices are responsible for the increased prevalence of nutritional deficiencies and disorders, infectious diseases, overweight and obesity, non-communicable diseases such as diabetes, hypertension, cardiovascular diseases, cancers and other chronic diseases. Sedentary lifestyles exacerbate the prevalence of nutrition and health problems. Conversely, dietary modifications to select and consume healthy diet have been proven to reduce diet related chronic diseases in the population.

The PDGN recommend exclusive breastfeeding to infants for the first six months of life followed by continuation of breast feeding at least till the age of two years along with appropriate complementary feeding from six months onwards.

The PDGN advise the daily consumption of basic food groups including cereals, meat and pulses, milk and milk products, vegetables and fruits. The guidelines suggest that the consumption of refined cereals should be reduced and substituted by wholegrain cereals while the consumption of fresh fruits, vegetables, pulses, meat and milk should be increased as per body requirements. It is further proposed that the amount of salt, sugar, saturated fatty acids, trans-fatty acids, ghee and cooking oil, sweets, soft drinks and juices containing added sugars, refined and processed foods should be limited.

The food handling and management practices should be safe, and the use of contaminated and untreated industrial wastewater must be prohibited. The pesticides, herbicides, antibiotics and the concentrations of heavy metals like lead, arsenic, mercury and cadmium in foods should be checked to remain within the

permissible range as recommended by the Codex Alimentarius Commission and Pakistan Standard Quality Control Authority, Pakistan.

In addition, regular exercise and physical activity are crucial for the prevention and management of diseases. The essence of PDGN is to provide updated dietary recommendations to general population, promote good nutrition, safe food handling and management practices and adopt healthy lifestyles for improved life quality and economic productivity.

The PDGN will be useful for multi-sectoral professionals and policy makers to formulate appropriate policy, plan and programmes. These guidelines will also facilitate Provincial Food and Drug Authorities to implement food quality standards in their respective provinces for ensuring that foods are safe and of better quality. The PDGN are intended to help in maintaining healthy body weight and reduce the incidence of under-nutrition, over-nutrition, infectious and non-communicable chronic diseases and premature deaths by promoting healthy diets.

INTRODUCTION

Optimal nutrition is essential for the physical and cognitive development of the body, behavioural and scholastic performance, better health and productivity, improved quality of life and socio-economic development of the country. Pakistan is challenged by food insecurity, poverty and by all forms of malnutrition, growing epidemic of non-communicable chronic and infectious diseases that have put an enormous burden on health services, resulting in avoidable disabilities and premature deaths.

A growing body of evidence suggests that unhealthy dietary and lifestyle practices are the key risk factors to the growing disease burden¹⁻⁴. Malnutrition accounts for half of the children's death, costs about 3% of yearly GDP in developing countries and results in losses of more than 10% of lifetime earning potentials in malnourished children⁵⁻⁶. Likewise increased and frequent consumption of energy dense nutrient poor foods with high salt, sugar and fats lead to over-weight and obesity and a consequent increase in the prevalence of metabolic syndromes, chronic diseases and premature deaths. It has been estimated that 340,000 cancer cases per year in the United States could be prevented with a healthy diet, combined with physical activity and maintaining a healthy weight⁷.

Nutrition transition and lifestyle changes have led to an increased prevalence of overweight and obesity and a pandemic of non-communicable diseases (NCDs) i.e., diabetes, cardiovascular disease, hypertension and cancer. NCDs are the leading causes of premature deaths in both the developed and developing countries accounting for 38 million of the world's 56 million deaths in 2012⁸. Expenditure on the treatment of NCDs amounts to billions of dollars annually adversely affecting the economy of countries, increasing disabilities, high premature deaths and impaired physical productivity and economic outputs. NCDs are responsible for loss of productive life years and cause innumerable sufferings for families and societies.

In Pakistan, an estimated 10-11% of the adult population (aged 25 years and above) is suffering from diabetes and the numbers are expected to increase to 14.5 million by the year 2025⁹⁻¹¹. In addition, cardiovascular diseases are one of the leading causes of disability and premature deaths in adults worldwide, low income to middle income countries bear the brunt by accounting for over 80% of the global disease burden and overwhelming health expenditures amounting to billions of dollars annually¹²⁻¹³. Similarly, CVDs account for 19% of all the deaths occurring in adults aged 30-70 years in the country¹⁴.

A synergistic relationship between nutrition and disease exists and good nutrition is essential for the prevention, control and treatment of diet related disease. Appropriate dietary practices and lifestyle modifications are required to prevent the rising prevalence of malnutrition, chronic diseases and premature deaths. Pakistan Dietary Guidelines for Better Nutrition (PDGN) are evidence-based dietary recommendations for maintaining optimum health and nutrition of the general population, which are derived on scientific knowledge about food consumption, dietary patterns, nutritional requirements, lifestyles and the relationship between diet and diseases.

PDGN provide a framework to guide the disease free normal population in the selection and consumption of a variety of safe and nutritious foods that are easily available, affordable, meet the daily nutritional requirements, boost the immune system against communicable and non-communicable diseases and

promote healthy lifestyles. Development of dietary guidelines require in depth analysis of population specific food consumption patterns, dietary habits, nutrients intake, nutrition and health status, socio-economic conditions and understanding of diet related public health issues of the population. In PDGN, emphasis has been laid on the consumption of a variety of safe and nutritious foods rather than nutrients, as the former is primarily consumed and easy to recall for dietary assessment and nutrients intake.

Rationale

- First national DGs were developed in 2005 by the Ministry of Health, Government of Pakistan to provide dietary recommendations for infants, children and adults. Lack of dietary diversification, unsatisfactory maternal and child care practices, industrialization and changing lifestyles in addition to natural catastrophes have led to a deteriorating nutrition situation in the country, that made it essential to review the relationship between diet and disease in the local context and revisit the existing food based dietary guidelines to mitigate the risk factors for under and over-nutrition and chronic diseases.
- The revised PDGN have taken into account the local dietary practices, cooking methods, cost of diet, nutrition and health situation of the population, socio-cultural practices, economic and environmental conditions to meet the nutritional requirements of individuals by and large.
- As food consumption and dietary patterns of individuals vary from country to country, so do the dietary guidelines developed to meet the nutritional requirements of the population on the basis of age, gender and physiological status. Similarly with the passage of time, there has been an increase in population growth, rural to urban migration as well as change in dietary habits, socio-economic conditions, lifestyles and prevalence of communicable and non-communicable diseases. Hence, dietary guidelines need to be revised to meet the changing population needs.
- The PDGN provide a list of foods with portion sizes to help the general public make smart and healthy food choices for a healthy, long and active life. Women of childbearing age, preschool children and adolescents are more vulnerable to nutritional deficiencies due to their increased physiological requirements and prevailing dietary and socio-cultural practices. The development and implementation of age specific dietary guidelines thus is vital for maintaining nutritional balance, weight management, prevention of diseases and improving the quality of life of the population particularly the most vulnerable groups.
- PDGN are resource for evidence-based decisions making and better policy choices. They also serve as tool for programme managers and professionals engaged in agriculture, food, nutrition and health related activities to develop cost-effective policies, strategies and nutrition programmes that promote healthy diet, support production, access and utilization of safe and nutritious foods, including interventions to reduce and control diet related diseases.

Goals

The goals for PDGN are to:

- Improve the nutritional wellbeing of the population by adopting and promoting healthy eating practices and lifestyles.
- Reduce the incidence of nutritional deficiencies and chronic diseases in the population.
- Improve the quality of life of the population and contribute to socio-economic development of the country.

Objectives

The objectives for PDGN are to:

- Use scientific evidence on dietary patterns in PDGN to provide basis for formulating relevant policies, strategies and plan of actions.
- Promote healthy dietary practices among the population to reduce and control nutritional deficiencies, diseases and deaths.
- Promote hygiene, safe food handling, proper preparation and improved storage practices to reduce and control food and water-borne diseases.
- Integrate physical activity into dietary interventions to achieve an effective and sustainable weight management programme.



NUTRITION SITUATION

Nutritional status of infants (0-6 months)

Early infancy (0-6 months) is a period of rapid body growth and brain development characterized by increased nutritional requirements that can only be ensured through exclusive breastfeeding. Exclusive breastfeeding is essential for optimum growth and development of infants during the first six months, and lays the foundation for future nutrition and health status of individuals. Mother's feed during the first six months is considered nutritionally complete and sufficient to meet the infant's nutrient and fluid requirements. Exclusive breastfeeding should be initiated within the first hour of birth and continued till six months of age without introducing any other food to the infant. Breast milk is rich in immunoglobulins and other anti-infective factors such as lysozymes, lacto-ferrin, interferon, anti-inflammatory and immune-modulating components that enhance immunity and reduce infant morbidity and mortality¹⁵. It has been reported that breastfed infants have at least six times greater chances of survival than non-breastfed infants¹⁶.



© FAO/Noomeena Anis

It has been reported that in Pakistan, only 24% of the women initiate early breastfeeding while exclusive breastfeeding under six months is estimated at 37% which falls under the category of low exclusive breastfeeding countries^{17,18}. The late initiation of breastfeeding and low prevalence of exclusively breastfeeding make infants more vulnerable to nutritional deficiencies and diseases. Promoting exclusive breastfeeding has a great potential to improve the nutritional and health status of the young children. Mothers and caregivers need to be encouraged to initiate breastfeeding within one hour of birth, exclusively breastfeed their babies for six months and continue breastfeeding along with appropriate complementary foods up to two years or beyond¹⁹. A longer duration of breastfeeding has been linked to reduced risk of childhood illnesses and obesity and improved cognitive outcomes.


Nutritional status of infants and young children (6-24 months)

Adequate infant and young child feeding is vital for growth and fundamental to the development of child's full human potential and for preventing chronic malnutrition which is manifested by high prevalence of stunting in Pakistan and other less developed countries. It is well recognized that the period between pregnancy and childhood up to two years of age (the first 1000 days of life) is a critical window for promoting optimal growth, health and behavioral development. It provides a window of opportunity to intervene, prevent growth faltering, malnutrition and diseases in children and provide a strong foundation for future physical and mental performance and improved quality of life. Nutrient requirements of infants after six

months of age increase which can't be met with breast milk alone, therefore, it must be complemented with good quality and quantity of safe and nutritious complementary foods. Breast feeding along with complementary feeding should be continued for at least two years and the amount of complementary foods should be increased gradually to ensure that foods are in better variety and quantity, sufficient, safe, nutritious and suited to the needs of the growing infant.

In Pakistan, poor IYCF practices are the leading cause of child morbidity and mortality. Poor IYCF in the form of delayed initiation of breastfeeding, less frequent and or inadequate breastfeeding, avoidance of giving colostrum, frequent use of pre-lacteals, use of breast milk substitutes, too early or too late introduction of quality complementary foods, inadequate postnatal care and poor hygienic practices make children vulnerable to nutritional deficiencies, infectious diseases and deaths. It has been reported that only 39% of infants receive complementary foods at the recommended time between 6-8 months²⁰.

Improved IYCF practices have a great potential to reduce malnutrition, diseases and deaths in children. At least 20% of child deaths could be prevented through improved infant and young child feeding practices, which are safe, non-invasive and cost effective as well¹⁵. Comprehensive, multi-sectoral integrated efforts are needed to protect, promote and support exclusive breastfeeding and complementary feeding practices essential for child survival. There is increasing recognition that optimal complementary feeding depends not only on what is fed, but also on how, when, where, and by whom the child is fed. It is essential to ensure that caregivers are provided with appropriate guidance regarding optimal feeding of infants and young children.



© FAO/Noameena Anis

Nutritional status of children (24-36 months)

Growth and development of children in this age group become slower than in the first two years of life along with improved digestive and enzymatic capacity, enhanced neural and hormonal responses which make children less vulnerable to nutritional deficiencies and diseases. Further, children in this age group are able to digest, absorb and utilize family foods and beverages and love to taste and eat family foods with minimum efforts of the parents. Children in this age group on an average gain about 2-2.5kg weight and grow 8-9 cm in height per annum²¹.

The slower growth and development of children elicits lower nutrients demand from exogenous sources and should be considered a period of “catch-up growth” to replenish body's nutrient stores and cover the earlier growth deficits. However, due to ignorance and lack of education, poor dietary practices, poor hygiene and sanitation, children in this age group are unable to cover their growth deficits and are increasingly susceptible to nutritional deficiencies and diseases. Rates of stunting, wasting and

underweight continue to rise in all the developing countries including Pakistan associated with a higher prevalence of diarrhea and acute respiratory infections which are the leading causes of morbidity and mortality in children²²⁻²⁴.

Nutritional status of children (3-10 years)

Growth and development during 3-10 years of age become relatively slower than the early childhood period but an adequate and well balanced diet is still required not only to cover the earlier age growth deficits but also meet the growth and development of the growing child. Children are relatively less vulnerable to nutritional deficiencies than infants and young children due to their improved digestive, physiological, metabolic, neurological and muscular coordination, slow growth and development and reduced nutritional requirements. Nature provides a better opportunity to children for catching-up on their earlier growth deficits, replenishing their body stores and making them ready for the adolescence growth spurt. During this period, children on an average gain about 2 kg weight and grow 5-6 cm in height per annum²¹. Pakistani children though show



improvement in their weight and height with increasing age but fail to recover significantly from early age nutritional deficits and their weight and height remain lower than the corresponding age reference children²². Similar results on anthropometric measurements of children were also reported by others^{25,26}.

Inadequate consumption of milk and milk products along with inadequate consumption of fortified foods in children's diet makes them prone to calcium, phosphorous and vitamin D deficiencies. These deficiencies are responsible for poor bone mass and are an increasing risk factor for rickets in children. Snacks consumption of children is limited to empty energy dense foods such as candies, sweets, toffees, soft drinks, samosas, papars, pakoras and rolls etc. that are unable to meet most of the macro and micronutrients requirements of the body²⁷. Similarly a large proportion of children don't regularly eat meat and meat products, fruits and vegetables either due to their unavailability at the household level or personal disliking and inadequate knowledge. Review of data on children dietary practices highlight that children's diets are lacking in nutrients dense foods, which increase the risk of nutritional deficiencies, malnutrition and diseases²⁸⁻²⁹.

Nutritional status of adolescents (10-19 years)

Adolescence is a period of transition between childhood and adulthood covering the age between 10-19 years³⁰. It is marked by a rapid increase in the reproductive, skeletal and physical growth and development of adolescents. The rapid physical and physiological growth provides the foundation for accomplishing present and future responsibilities more efficiently that can only be ensured with adequate and optimal nutrition. Conversely, lack of a balanced, nutritionally optimal diet at this critical juncture puts individuals and their future generations at a greater risk of nutritional deficiencies and makes them more vulnerable to diseases³¹.

Adolescence exerts greater nutritional demands to meet an increasing body needs for nutrients and is greatly influenced by poor food choices and unhealthy eating habits. Eating disorders are more common during this crucial transitional stage of growth and can be attributed to a variety of factors such as peer pressure, body image consciousness, lack of self-esteem and other social influences. Adolescents with severe eating disorders suffer from impaired physical and psychological growth and development.

Review of data on Pakistani adolescents nutritional status revealed that about one-third of the adolescents are either stunted or underweight and around 10-30% are overweight and obese, however, the prevalence of under and over-nutrition in adolescents varies from place to place due to differences in dietary patterns, socio-economic conditions and other environmental factors^{32,33}.

Dietary practices of adolescents show that skipping breakfast is more prevalent in girls than boys and there is no significant difference in the frequencies of skipping breakfast between the different socio-economic groups²⁷. Tea with bread and curry are the more commonly consumed breakfast foods of adolescents of lower socio-economic group while milk, juice, fruit, cereals, paratha and eggs were more often consumed in breakfast by higher income group. Candies, samosas, papars, pakoras, rolls, potato chips, grams and beans are largely consumed as snacks by adolescents of lower socio-economic groups while ice cream, fruit chat, burger, French fries and soft drinks are mostly consumed as snacks by the higher income groups²⁷.

The frequency of dairy, fruits and meat consumption indicates that at least two-thirds of the adolescents don't meet the recommended amounts and are therefore, more likely to be deficient in meeting the recommended daily dietary intakes of iron, zinc, calcium, phosphorus, vitamin D and many other nutrients.



© FAO/Noomeena Anis

Nutritional status of adults (19-60 years)

Adults in Pakistan are also confronted with the problems of under and over nutrition with an alarmingly high prevalence of non-communicable diseases. About a quarter or more of adults are deficient in one or more nutrients like iron, vitamin A, calcium, vitamin D, zinc and iodine and are suffering from diabetes, hypertension, cardiovascular diseases and cancer¹. Cardiovascular diseases (CVDs) are one of the leading causes of disability and premature deaths in adults worldwide. Low to middle income countries bear the brunt by accounting for over 80% of the global disease burden and overwhelming health expenditures amounting to billions of dollars annually³⁴. Women in Pakistan are more vulnerable to nutritional deficiencies and non-communicable diseases than men with 18% underweight, 19% overweight, 10% obese, 50% anemic, 42% vitamin A and zinc deficient that are attributed to poor dietary practices, lower access to health care services and heavier workloads with multiple domestic and outside jobs^{1,35,36}. It has been reported that women with a mean age of 52 years had a higher prevalence of coronary artery disease i.e., 30% than that of 23.7% for corresponding age men³⁶.

The National Health Survey (1990-94) report revealed that 1 of 3 individuals aged ≥ 45 years suffer from hypertension³⁶ while another study reported that 22% of adult population was suffering from diabetes³⁵. The prevalence of diabetes was higher in men i.e., 23.7% as compared to 20.6% for women³⁵. Statistics on demographic, economic, social, nutrition and health indicators of Pakistan revealed that though there is improvement in per capita income³⁷ and household possession of electric goods, transportation and communication facilities, yet no substantial improvement in the nutrition and health indicators of the country. For better health outcomes and improved quality of life, concrete measures are needed to address the nutrition and public health issues.

Nutritional status of elderly (>60 years)

In Pakistan, the older population constitutes 4% of the total population and the proportion is likely to increase over the period of time³⁸. In 2018, the elderly population is projected 6% (PBS). The advancement of medical technologies and diagnostic facilities and a surge in per capita income has led to an increase in the life expectancy and the proportion of older population. The elderly because of their waning health and compromised physiological functions i.e., decreased mastication, salivation, gastric secretion, digestive and absorptive capacity combined with decreased physical activity and appetite, loss of lean body mass and skeletal muscle, need special diet and health care to reduce the risk of physical and mental disabilities and control of diseases.

Looking after the health and social needs of older people is becoming a challenge both for the poor families and the government due to less sensitization of the issue, limited commitment and inadequate resource mobilization by the government to provide dietary, health, social and other related services particularly to the poor families in meeting their nutritional requirements.

It has been estimated that in Pakistan, over 72% of the older people have five or more health problems but there has been hardly any nutrition and health care programme to address their nutrition and health needs³⁸. Diabetes, hypertension, coronary artery disease, arthritis and cancer are on the rise, causing an

increase in the family sufferings due to high medical costs, increased care, leading to overburden on the health care system with losses of billions of rupees and increased early deaths of elderly. Financially sound older people get relatively better nutrition and health care services but older people belonging to poor families are the ones who are under tremendous pressure, stress and are mostly unable to get proper nutrition and health care, which results in increasing burden on both the families and government. In Pakistan, limited or negligible work has been done on the older population to assess their nutrients requirements and intake, however, small cross-sectional studies revealed that the diet of the older people is deficient in almost all the macro and micronutrients^{39,40}.

The existing food consumption data on older population revealed that their diet is mainly energy dense and less diversified, which make them vulnerable to nutritional deficiencies with impaired immune functions and increased risk to acute and chronic diseases. Comprehensive, holistic, cost-effective dietary and health care interventions including nutritional assessment, counseling, dietary supplementation, and establishment of nursing homes for older people with improved health and social care are needed to enable them to participate in the socio-economic development of the country, reduce their sufferings and improve their quality of lives.

Nutritional status of pregnant women

Pregnant women are more vulnerable to nutritional deficiencies and health problems due to increased nutritional requirements to meet their physiological needs, sustain foetal growth and development. Decreased nutrients intake during pregnancy has been attributed to cultural, social and economic reasons. In Pakistan, early marriages coupled with poor dietary intakes and lack of proper antenatal health care further compromise the nutritional status of pregnant women with an increased prevalence of nutritional deficiencies.

According to NNS 2011, 62% of the pregnant women sought antenatal care (ANC) being greater in urban (81.4%) than rural settings (53.7%)¹. The low utilization of ANC in rural areas could be due to lack of awareness, unavailability or long distance from home to ANC facilities and socio-cultural reasons that may restrict women mobility. Similarly it had been reported that only a quarter of women received vitamin-iron supplements during pregnancy. The low consumption of supplements by women was attributed to lack of knowledge about the need and importance of supplements and food sources of vitamin-minerals¹.

Poor dietary practices among pregnant women were reflected by the increased prevalence of malnutrition i.e., about 50% of the women were either underweight, overweight or obese, 16% had night blindness, 51% were anaemic, 37% iron deficient, 46% vitamin A deficient, 48% zinc deficient, 69% vitamin D deficient and 59% were calcium deficient¹. These results were substantiated by others⁴¹ who reported that only 57.3% and 46% of pregnant women increased their food frequency as well as food intakes, respectively, 61% took dietary supplements where as 17.3% made no changes in their dietary intakes. A good proportion of women reported food cravings 78%, aversion 38% and avoidance to certain food 22%.

A dietary intake study on Pakistani-Norwegian (PN) pregnant women revealed that PN women had a significantly lower intake of vitamin D (2.2 vs 3.3 µg/day), calcium (793 vs 1134 mg/day), nutrition

supplements (2.9 vs 7.0 µg/day) and lower serum vitamin D levels (19 nmol/l vs 55 nmol/l) than that of the Norwegian women. In addition, 83% of the PN women were found vitamin D deficient having serum 25-hydroxyvitamin D3 levels below the reference value (< 30 nmol/l)⁴².

Similar results on dietary intake of pregnant women were reported by others⁴³ who observed that regardless of the trimester and nutritional status, pregnant women were consuming low calories (mean 1437 Kcal/day in the first trimester; 1407 Kcal/day in the third trimester). The study further revealed that women were consuming less than the recommended amount of various foods i.e., 56% were consuming less than the recommended servings of bread and cereals, 80% less than the recommended servings of meat, 94% less than the recommended servings of milk, 96% and 94% less than the recommended servings of fruits and vegetables, respectively. In addition, undernourished women had significantly lower body mass indices than those of the well-nourished women both in the first and third trimesters.

The dietary intake and nutritional status of pregnant women were further substantiated by the fact that 22% of women were underweight⁴⁴. Only about 49.61% of the adult non-pregnant and non-lactating women, 46.92% of pregnant women and 32.31% lactating women had recommended caloric intakes⁴⁴. Other studies conducted on Pakistani pregnant women revealed similar results of dietary inadequacy and increased risk of adverse pregnancy outcomes⁴⁵⁻⁴⁷. Keeping in view the deplorable nutritional situation of pregnant women, comprehensive integrated nutrition and health care interventions are required to improve the nutritional and health status of pregnant women for desirable pregnancy outcomes and averting tragic consequences in the form of stillbirths, neonatal, infant and maternal mortalities.

Nutritional status of lactating women

Lactating women have increased nutritional requirements due to their body's increased physiological needs for breast feeding the baby. On an average, lactating women in the first six months of lactation produce 750-800 ml of milk per day which provides approximately 525-560 Kcal/day to the infants⁴⁸.

It has been agreed that breast milk is best for infants in terms of its ideal composition, qualitatively and quantitatively sufficient for the first six months of infants, confers immunity and reduces the risk of diarrheal diseases by four to fourteen folds and the risk of respiratory infections by five folds⁴⁹. There is also general consensus that healthy lactating women from developing countries produce more or less the same amount of breast milk comparable to lactating women of developed countries but maternal nutritional status has varying effects on nutrients composition of breast milk^{50,51}.

Studies have shown the association between healthy diet and quality of breast milk. Studies revealed that supply of nutrients like vitamin B1, B2, B6, B12, choline, retinol, vitamin A, vitamin D, selenium, iodine and folate to the baby is decreased by maternal nutrients deficiencies while it has no effect on calcium, iron, copper and zinc supply to infants⁵¹. For adequate milk output and supply to the infant, good maternal nutritional and health status, appropriate management of lactation, avoidance of pre-lacteals and formula feeding is crucial.

Due to ignorance, lactating women pay very little attention to their diet which adversely affects the


nutritional status of both the women and baby as depicted by the high prevalence of malnutrition in women and children^{1,43}. Focused and targeted nutrition interventions to improve lactation management and dietary diversity with emphasis on increased quality and quantity of macro and micronutrients are essential for the proper growth and development of infants and children.

Food availability and consumption in Pakistan

In Pakistan, food availability (kg/capita/year) has remained relatively stable over the last three decades to meet the demand of growing population. The availability of major food items i.e., wheat, rice, maize, fats and oils, meat, milk, vegetables, fruits and pulses which indicates that by and large food availability has improved over the years and as such there had been no shortage or disappearance of food from markets (Figures 1-4)^{52,53}.




Figure 1. Availability of cereals over different time periods




Source: MPDR, 2016


Figure 2. Availability of fats and cooking oils over different time periods



Source: FAO, 2016

Figure 3. Availability of maize and sugar over different time periods

Source: FAO, 2016


Figure 4. Availability of pulses, fruits and vegetables over different time periods

Source: MPDR, 2016

Based on the food availability data, the estimated per capita calories availability were 2470 calories per day⁵⁴. The availability of milk, meat, sugar, fats and oil has markedly increased over the years. The food availability data demonstrate that Pakistan does not fall under the category of food deficit countries and

local production is sufficient to meet the food requirements of the population with an exception of a few food items which are imported to bridge the demand. At the same time, an appreciable amount of rice, wheat, sugar, fruits and marine foods are exported to earn foreign exchange. The country, however, is lagging behind in the export of value added foods and there is lot of potential for value addition and export of food commodities. Although the income generated has increased over the years yet is insufficient for the common man to purchase the diversified foods and meet the recommended nutrients requirement of the body (Figure 5)³⁷.


Figure 5. Per capita GDP from 1960-2015



Source: World Bank, 2016

Household income has been a major constraint for purchasing and consuming quality foods for majority of the population. It has been estimated that using cost for basic needs method, 29.5% of the population live below the poverty line⁵⁵ while caloric based poverty assessment revealed that 35% of the population is unable to purchase foods sufficient enough to meet the nutritional requirements of the body⁵⁶ (Figure 6).

Figure 6. Poverty based on low calories consumption over the years



Source: Nazli et al., 2014

Imbalance between body's nutrient requirements and dietary intake results in malnutrition, which is affecting a large segment of the population impeding their work potential and earnings. A significant proportion of Pakistani population has a deficiency of both macro and micronutrients on the basis of an established notion that a diet deficient in meeting the energy needs of individuals would also be deficient in supplying the recommended nutrients intakes for all essential nutrients⁵⁷. In other words, energy could be used as a marker of individual nutrients intake i.e., reduced energy intake would result in a deficient supply of all other nutrients.

FAO has produced estimates of the Prevalence of Undernourishment (PoU) and of the Number of Undernourished (NoU) since 1974. In addition, the country level estimates of the PoU and NoU, in addition to regional and global aggregates are published in the State of Food Insecurity report (SoFI). The methodology for estimating the PoU is based on the comparison of a probability distribution of habitual daily Dietary Energy Consumption and a threshold level, called the Minimum Dietary Energy Requirement (MDER)⁵⁸. In Pakistan, proportion of undernourished in total population is 22% as per State of Food Insecurity report 2015⁵⁹.


Relationship between diet and disease

Diet has been recognized as one of the strongest predictors of health and disease but its role and importance has largely been underestimated due to unawareness, lack of knowledge, preference for traditional monotonous diets driven by economic and cultural reasons. Studies conducted around the world revealed that unhealthy dietary habits and physical inactivity are the major determinants of overweight, obesity and chronic diseases and are responsible for decreased productivity and increased mortality⁶⁰.

Excess salt intake has been linked to 1.7 million deaths from cardiovascular diseases and it has been reported that a small reduction in salt intake was associated with more than 7% reduction in deaths from strokes and heart attacks⁶¹. Similarly, it has been estimated that at least one third of cancer cases can be prevented by adopting healthy dietary practices and improved lifestyles⁶². Changes in dietary habits and lifestyles can make a substantial reduction in diet related chronic diseases and improvements in nutritional and health status of the population.

Data on food consumption patterns revealed that diet of both poor and non-poor households is predominantly comprised of energy dense foods with little diversity and therefore micronutrient deficiencies are more likely to be prevalent in the population unless foods are fortified (Figure 7)⁶³. The figure indicates that wheat is the main staple providing 46 to 52% of the total daily caloric intake, oils 14 to 13%, dairy 13-10%, sugars 10% and meat 3 to 2% of the total calories for the non-poor to poor groups of population, respectively.


Figure 7. Calories contribution of different foods in poor and non-poor diets



Source: Malik et al., 2015


Figures 8-9 reveal that the dietary patterns of the rural population remained almost similar to that of the urban populations except that the former had a higher consumption of milk than the later which could be attributed to greater accessibility of the rural population to milk from buffaloes, cows, goats and sheep that are mostly reared in the rural areas. The figures also showed that diet of the population was less diversified and therefore, less likely to meet all the nutrients requirement of the body

Figure 8. Food consumption patterns of households.



Source: Malik et al., 2015


Figure 9. Frequency of food consumption based on 7 days dietary recall



Source: Nazli et al., 2014

Accurate assessment of food consumption, nutrients intake and requirement by the population is crucial to formulate food based dietary guidelines to prevent the incidence of malnutrition and chronic diseases in the population. However, it is important that food should be available, accessible and affordable at all times so that a variety of foods can be purchased to supply adequate amounts of nutrients to the body. The estimate suggests that households with low income spend over 59% of their income on energy dense foods and hardly any amount is left for education and health care of the family⁶³. Besides, there has been a trend of increasing food prices which limit the purchasing power of the common man to purchase and consume nutrient dense foods that exacerbate food insecurity and malnutrition (Figure10).

Figure 10 . Trends in food prices over the years



Source: Malik et al., 2015

It is well established that unhealthy dietary practices and physical inactivity are risk factors for overweight and obesity and are associated with impaired metabolic functions, inflammation and increased prevalence of non-communicable disease. Food consumption patterns (FCPs) in Pakistan reveal that majority of the population consume refined cereals, hydrogenated and un-hydrogenated oils with high consumption of salt and sugars which are far above the recommended nutrients intake and dietary recommendations of World Health Organization⁶⁴. The FCPs also show that consumption of vegetables, fruits, milk and milk products, meat and meat products is much lower than the recommended intake. Increased urbanization, economic and nutrition transitions have led to increased consumption of fast and processed foods, sweets and soft drinks with high energy, fat, salt and sugar content resulting in increased risks of malnutrition and chronic diseases in the country. Lack of appropriate policies and non-enforcement of food safety regulations have multiplied the number of street food vendors who sell low quality foods in schools, markets, canteens and other public places, increasing the prevalence of food borne diseases, malnutrition and mortality.

FCPs demand that dietary modifications may be encouraged transforming from energy dense foods to nutrients dense diversified foods that encompass a variety of food groups to meet all the essential nutrients to combat the double burden of malnutrition and chronic diseases. It requires multidimensional interventions and approaches to facilitate the dietary transition from energy dense to nutrients dense foods to control and reduce nutritional deficiencies and associated diseases. Furthermore, increased investments in social sectors are required to provide safe water, food and health care services as well as create more job opportunities for marginalized population to improve their quality of life.

Traditional and fast foods

In Pakistan, fast foods may be defined ready to eat food available in public places, food streets, food corners and restaurants. They may include burgers, shami kababs, pizzas, sandwiches, chat, cholay, samosas, pakoras, haleem, french fries, ice creams, doughnuts and all other traditional and non-traditional foods that are dispensed quickly. They are classified as energy dense foods because they provide high amounts of sodium, sugar, cholesterol and fat but are low in vitamins and minerals and dietary fibre content. Despite of their low nutritional value, fast foods have become a symbol of modern culture and their taste appeal more to children and youth.

Likewise other countries, consumption of fast food in Pakistan is also flourishing. According to various reports, fast food industry is the 2nd largest industry in Pakistan; accounting to 27% of its value-added production and 16% of the total employment in the manufacturing sector with an estimated 169 million consumers, Pakistan holds the world's eighth largest market when it comes to fast food and food related business. More than 1000 large-scale food processing enterprises in Pakistan are in operation⁶⁵.

Traditional foods are slowly losing their appeal, as fast foods are alluring to the population due to their convenience and easy availability. Home cooking hours have been comparatively minimized and people prefer eating outside in food streets and fast food restaurants. Along with Pakistani foods, Chinese, Western, Italian and Continental cuisines have also gained a lot of liking and are becoming a part of the Pakistani menu.

The impact of fast foods on health is quite staggering when consumed on a regular basis. By replacing fresh wholesome food with processed additive-laden food, it has been noted that blood pressure rises, cholesterol level increases, metabolic rate altered and the immune system is weakened.

Glycemic index of foods

It is the ability of carbohydrate containing foods to raise the body's blood glucose levels after their consumption. Foods by virtue of variations in the quality and quantity of carbohydrate contents yield different glycemic indices upon their consumption. In general, refined cereals/grains (white bread, white rice, spaghetti and processed foods) have a higher glycemic index than whole cereals/grains (whole wheat bread, brown rice). Foods are classified as having high (GI 70 or more), medium (GI 56-69) and low (GI 55 or less) glycemic indices⁶⁶. Glycemic load is a multiple of carbohydrate quality and quantity that tends to measure the effect of food on postprandial blood glucose level. Foods with high glycemic indices put greater stress on the body to secrete greater amount of insulin to maintain the rising blood glucose level to a normal blood glucose level. In addition, higher carbohydrates intake adversely affects glucose and lipid metabolism leading to an increasing risk of cardiovascular diseases⁶⁷. Conversely, low glycemic index diets have been associated to improving the overall blood glucose control in people with diabetes⁶⁸. A list of foods containing different glycemic indices is given as Annexure-I.

Food allergy

The abnormal response or reaction triggered by the body's immune system to certain foods is called food allergy. Foods that cause allergy are called allergen. The allergic reaction may be of varying intensity from mild to severe, may affect any part of the body and could be for a shorter or longer time period. Skin, gastrointestinal tract, cardiovascular system and respiratory tract are more commonly affected. Food allergy affects about 4-6% of children and adults⁶⁹.

The common symptoms include skin irritation and rash, itchy sensation in mouth, swelling of the lips, tongue or throat, shortness of breath, trouble breathing, flushing, wheezing, stomach pain, vomiting and diarrhoea etc. The reactions are mediated by immunoglobulin E and other antibodies. Foods that often trigger allergic reactions are milk, eggs, peanuts, nuts, fish, soya, wheat and shellfish (shrimp)⁷⁰.

Unprocessed and processed foods

Pakistan is a small but growing market for imported consumer food products and the small modern retail sector is growing slowly. Rising incomes, urbanization, and a young populace are combining to slowly shift traditional consumption patterns away from bulk and raw foods towards packaged and processed foods, including ready-to-eat meals and frozen foods.

A typical Pakistani household makes regular purchases of staple foods (i.e., wheat flour, pulses, edible oils, fruits, vegetables, milk, and meat) several times per month from neighborhood stores due to convenience, perceived freshness, and limited storage space in home. Young Pakistani professionals (men and women)

are making monthly food purchases from modern retail stores due to greater variety of available products, access to restaurant/food corners and prepared foods all under one roof. For urban dwellers with a sufficient income, one-stop grocery shopping is catching on as a family outing.

Affluent Pakistani families are attracted to modern retail stores due to their affordability, strategic locations, variety, and access to imported processed foods. The demand for specialty and high value foods such as dates, cereals, beverages, chocolates, almonds, cakes, fruits and fruit juices reaches its peak during the Islamic festive season, especially at Ramadan and Eid. These festivals revolve around the year depending on the lunar calendar. Fresh foods, fruit juices, fruit concentrate based beverages, organic foods, sugar-free confectionaries, packaged food with higher fiber content, dairy products, and vitamin and calcium fortified packaged foods and beverages are all gaining acceptance among high income consumers.

Traditional and modern snack foods, such as confectionaries, pastries, cakes, biscuits, ice cream, or sweet and savory snacks are very popular among Pakistanis. Frozen foods and instant noodles, which are easy to prepare for children, are popular among working mothers. Local flavors are preferred and local food manufacturers are exploring opportunities to produce new products using a combination of local and imported flavors. The rise of the urban middle class has increased the acceptance of packaged and ready-to-eat food products. Many Pakistanis are quite willing to try new foods while eating out, but often return to traditional foods at home. Chinese, Italian, Thai, Moroccan, and Lebanese foods are among the fastest growing new cuisines in Pakistan.

Multinational companies have also been successful in capturing the local food markets in Pakistan, which are encouraging youth to go for processed foods. Frequent consumption of processed foods is injurious to health due to the excessive use of salt, free sugars and other additives to increase the shelf life of the products. While Pakistan is slowly switching to processed foods, a reverse trend is being noticed in the Western world with many countries promoting a greater use of organic and homegrown natural foods. More and more consumers are buying fresh produce from the many farmers' markets that have sprung up in the urban areas.

To fight the epidemic of obesity, consumers are constantly reminded to reach out for fresh fruits and vegetables while avoiding sugary drinks and fried items. Making healthy food choices is vital to control this growing epidemic. Processed foods are treated with multiple chemicals to give them colour, taste, longevity and freshness and thus are exposed to energy-intensive processing. In addition to this, cardboard packaging often contains chemicals called diisopropyl-naphthalenes (DIPNs), which can migrate into the food contents rendering it unhealthy for consumption⁷¹.

Nutrition labeling

Nutrition labeling is the information printed on the packages by food manufacturers to inform consumers about the name and total amount of food, list of ingredients, its nutrients contents, the percentage of daily nutrients reference values per 100 g, per 100 ml or package of the food. In developed countries it is mandatory for the food industry to label all pre-packaged foods with their nutrients composition that are marketed for human consumption. Nutrition labeling assists consumers in selection of foods according to their taste, liking and nutritional needs. It enables consumers to make wise choices and take informed

dietary decisions. It provides information on the amount of nutrients such as sugar, salt, saturated fatty acids, cholesterol, cis and trans fatty acids, the increased levels of which are considered harmful to health. The information on nutrition labeling varies from country to country determined by the respective food regulatory bodies.

Overweight and obesity

Energy intake greater than body requirements leads to an increased risk of overweight and obesity, which are strongly linked with increased incidence of diabetes, hypertension, coronary artery disease, cancers and many other chronic non-communicable diseases. In Pakistan, consumption of energy dense foods (sweets, bakery products, fried foods, fast foods and overuse of oils and ghee in cooking) in combination with sedentary life styles are the main causes of overweight and obesity. It has been estimated that about 1/3rd of the adult population is overweight and obese¹. Weight management through diet and lifestyle modifications is most effective in preventing and treating overweight and obesity.

Body Mass Index (BMI) is a measure of body's nutritional status assessed by taking weight of adults in kilograms divided by height in meters square. WHO has proposed the following classification to categorize adults into underweight, normal (weight), overweight and obese⁷².

| Group | BMI |
|------------------------|----------------|
| Severely thin | <16.0 |
| Moderately thin | 16.0-16.9 |
| Mildly thin | 17.0-18.4 |
| Underweight | Less than 18.5 |
| Normal | 18.5-24.9 |
| Overweight | >25 |
| Pre-obese | 25.0-29.9 |
| Obese Class I | 30.0-34.9 |
| Obese Class II | 35.0-39.9 |
| Obese Class III | ≥ 40.0 |

Measurement of waist circumference is another useful indicator of assessing abdominal obesity. Excess abdominal fat is an independent risk factor for chronic diseases. Men having a waist circumference greater than 94 centimeters and women having a waist circumference greater than 80 centimeters are at a greater risk of metabolic syndromes, diabetes, hypertension and cardiovascular diseases⁷³.

Caloric reduction and increase in physical activity are two well-established and successful interventions to reduce and control overweight and obesity. Reducing caloric intake without affecting and compromising other nutrients intake would be a useful strategy for weight management. An individual's nutritional assessment is required for prescribing any dietary regimen; however, for prevention of overweight and obesity, it is important that an individual should follow PDGN on a regular basis. For treatment of

effective in reducing biomarkers of inflammation and better management of cancers, diabetes and cardiovascular diseases⁷⁴.

Physical activity and lifestyle behaviour

Physical activity is vital for weight management, bone and muscle strength, optimal physiological functions, emotional wellbeing and good health. Sedentary lifestyles promote overweight and obesity, impair metabolic functions and increase the risk of chronic diseases. Physical inactivity is the fourth leading risk factors for global mortality and has been associated with non-communicable diseases, metabolic disorders, and mental diseases⁷⁵. Evidence shows that engagement in optimum levels of physical activity improves cardio-respiratory and muscular fitness, bone health, metabolic functions and mental health. Weight reduction reduces health risks and helps in better management of chronic diseases⁷⁶.



Regular physical activity is one of the major contributors to wellness, quality of life and disease prevention. Moderate physical activity can lower the risk of heart disease, stroke, diabetes, osteoporosis, dementia, and numerous cancers. Though there are immediate benefits of physical activity, the majority of prevention and health benefits are achieved when it is practiced lifelong⁷⁷.

Economic and nutrition transition in both developed and less developed countries have caused substantial changes in dietary patterns and lifestyles of the people from complex unrefined foods to more refined and processed foods, manual physical work has been replaced by mechanical work and outdoor games are replaced by indoor sedentary computer games. These changes have been responsible for the pandemic of metabolic syndromes and chronic diseases affecting millions of people around the world with impaired physical and cognitive functions, increased morbidity, heavy burden on the health care systems and premature deaths⁷⁸.

The development of healthy eating behaviour and physical activity patterns helps to optimize health status and promote mental and physical wellbeing. Unfortunately, the population by and large is actively engaged in unhealthy eating behaviours such as frequent dieting, meal skipping and frequent consumption of foods high in total and saturated fats, sodium and sugar^{27, 79}. Physical inactivity is also very common particularly among adolescents.

In order to meet the challenge of improving physical activity and promoting healthy dietary habits, integrated efforts of parents, educators, health care providers, schools, communities, the food industry, policymakers and the people themselves need to be strengthened, all working together to create more opportunities for healthful eating and healthy lifestyles⁷⁵.

Promotion of healthy lifestyles in children focuses predominantly on proper nutrition and physical activity,

elements now widely recognized as essential for a healthy life. Systematic reviews have shown that healthy lifestyle such as increased levels of physical activity, a healthy diet, reduced stress and better sleep enhance general well-being⁸⁰.

There is scant data on physical activity in the Pakistani population. Moreover, the tools for assessing physical activity have not been validated in the Pakistani setting; it is important for future epidemiological surveys that such tools be developed, standardized and validated. Physical activity offers additional advantages for risk reduction in the Pakistani population since the highly prevalent risk factors amongst this population such as low HDL, central obesity and insulin resistance respond better to exercise. Stepping up physical activity in a population necessitates bringing a cultural change. This has two-fold implications; the first involving modification of lifestyles and the second necessitating the creation of a physical and social environment conducive to physical activity.

Smoking and smokeless tobacco use

Smoking has long been implicated as a risk factor for many chronic diseases, including cardiovascular, respiratory and gastrointestinal diseases and a variety of cancers. Tobacco smoke contains many oxidants and free radicals that can cause damage to lipids, proteins, DNA, carbohydrates and other bio-molecules. The annual death toll attributable to tobacco use is about 6 million while more than 600,000 are non-smokers being exposed to second-hand smoke⁸¹. In addition it has been estimated that 80% of 1 billion smokers worldwide live in low to middle income countries, the devastation caused by tobacco goes much beyond this picture with implications for individuals, societies and the health systems in general. In Pakistan, it has been reported that 45% of men and 6% of women smoke some type of tobacco⁸² which are lower than previously reported studies that indicated 54% men and 20% women using tobacco in one form or the other; prevalence of smoking increased with age among both men and women. Men in the age range of 25-44 years had the highest prevalence of smoking (cigarettes and beedis) whereas with respect to women, prevalence was highest between the ages of 45-64 years⁹. It was also reported that smoking was more prevalent in rural than urban areas and illiterate than literate populations⁸².

Cigarette consumption constitutes the single most important modifiable risk factor for coronary artery disease (CAD) and is the leading preventable cause of early deaths. Smoking has a particularly staggering impact in the less developed countries, almost half a billion individuals worldwide will eventually die of smoking-related complications⁹. Even among non-smokers, it has been recognized that inhaled smoke, whether from passive exposure or from cigar or pipe consumption greatly increase the risk of CAD.

Smoking alone is responsible for 10% of cardiovascular diseases⁸³ and those who consume 20 or more cigarettes daily have a two to three fold higher risk of CAD⁹. Moreover, these effects depend on the dose; consumption of as few as one to four cigarettes daily increases the risk of CAD. Smoking acts synergistically with oral contraceptive agents, placing younger women at an even higher relative risk. In addition to myocardial infarction, cigarette consumption directly relates to increased rates of sudden death, aortic aneurysm formation, symptomatic peripheral vascular disease and ischemic stroke. As for coronary artery disease, the risk of stroke is directly increased with the number of cigarettes consumed.

Food safety and hygiene practices

Food safety is an important pillar of food security. It ensures the availability of food that is safe, free from infectious agents and fit for human consumption. Access to safe foods and safe drinking water is essential for preventing infectious and chronic diseases. Microbiological and chemical contamination is a major cause of food and water borne diseases. Unsafe food contains harmful bacteria, viruses, parasites and chemical substances causing more than 200 diseases ranging from diarrhea to cancers⁸⁴. It has been reported that contaminated food and water affects millions of people around the world resulting in 4 million deaths in which developing countries take a major share^{85,86}.

Food hygiene is not given due importance neither at the household level nor at the commercial (restaurant, food streets and other outdoor eating places) level. Personal and food hygiene (equipment, utensils, food handling and kitchen surfaces) practices before, during and after cooking are far below the desired quality standards. Vegetables, fruits, meats and other raw foods are not properly washed, cleaned and stored. Water used for cooking is not clean and safe. Poor food storage systems in homes and restaurants increase people's vulnerability to frequent food-borne diseases and increase mortality. In addition, at the national level there are no food safety policies, strategies and work plans to implement and protect public health from consumption of adulterated and contaminated foods.

Hygienic food management

Promoting good personal and food hygiene practices at all levels of food handling starting from farm to table i.e., production, transportation, processing, storage and marketing are not only crucial for ensuring food security but also for human safety and survival. All food industries, cafeterias, restaurants, food streets and food vendors should be bound to follow good hygiene and food safety practices during food handling, cooking and servings to the public. Food safety rules and regulations pertaining to prohibition of exposing foods and beverages in the open air and sun and not properly covering or protecting them, shall be included and implemented strictly at all levels of food and beverages production and marketing. Rigorous monitoring and implementation of food safety laws at all levels of food chain, are fundamentally important for ensuring a regular supply of safe foods and clean drinking water to the public.

Sensitization and capacity building of officials from the ministries and departments related to agriculture, food, health, education and industry, food vendors and workers involved in food industry are essential for promoting good hygiene practices and providing safe food to the public. Food industry shall also be bound to label all foods and beverages produced in the country and a stringent policy of collecting and analyzing food and water samples for pathogens, adulterants and contaminants shall be adopted at all levels of production and marketing. Public awareness to adopt and promote good hygiene practices through electronic, print and social media would be useful and effective means of educating people to adopt good hygiene practices. Good personal and food hygiene guidelines have been proposed in Annexure-II.

Water, sanitation and hygiene

Water is a basic necessity of life, which must be clean and safe for human consumption. It should be free from pollutants, industrial wastes, physical, chemical and microbial contaminants that are injurious to

human health. Consumption of water that is contaminated with animal and human excreta is a major source of pathogenic bacteria, viruses, protozoa and helminthes and results in epidemics of waterborne diseases. The organisms most commonly found in water contaminated with sewage are Escherichia coli and the coliform group as a whole⁸⁶.

It has been reported that in Pakistan 20 to 40% beds in hospitals are occupied by patients suffering from water related diseases especially during the summer months⁸⁷. Unfortunately, most of the water sources are contaminated and there is no proper system to ensure the supply of clean drinking water to the marginalized populations. Estimates also suggest that in Pakistan a quarter of all the people visiting hospitals are suffering from water-related diseases as a result of poor sanitation and lack of safe drinking water facilities⁸⁸.

Countries providing clean water and sanitation have greatly reduced child morbidity and mortality. The environmental pollution in Pakistan has exacerbated the prevalence of chronic diseases and increased mortality⁸⁹. Improper waste disposal system, lack of safe drinking water, and lack of awareness about health and nutrition are the major concerns. The use of contaminated water and exposure to a polluted environment increase the risk of diseases such as diarrhea, dysentery, malaria, respiratory infections, influenza, and hepatitis. It is estimated that 45% of the total population in Pakistan does not have access to health services, 40% are deprived of safe drinking water and 53% are living without sanitation facilities⁸⁹. Drinking water is considered safe when it is free from microorganisms as well as from toxic chemicals of industrial wastes and heavy metals that are injurious to health. Avoid drinking untreated, surface and ground water that may be contaminated with industrial wastes, chemicals and microorganisms. Water at the point of origin may be treated by antimicrobial and anti-contaminant agents. At home drinking water can be made safe by boiling or chlorination or installing membrane filters or by thermal treatment. Chlorination at a concentration between 0.2-0.5 mg/L from normal to high risks circumstances is recommended⁹⁰.

The maximum allowable concentration of lead, selenium, arsenic, chromium and cyanide in drinking water are 0.1 mg/L, 0.05 mg/L, 0.2 mg/L, 0.05 mg/L and 0.01 mg/L respectively, drinking water containing metal concentrations above these limits shall be considered unfit for human consumption⁸⁶.

METHODOLOGY

Under the Scaling up Nutrition (SUN) initiative, the Nutrition Section, Ministry of Planning, Development and Reform (MPDR)/ Planning Commission and Food and Agriculture Organization of the United Nations (FAO) joined hands in formulation of Pakistan Dietary Guidelines for Better Nutrition (PDGN). A brainstorming workshop was held in which renowned experts from the Ministries, universities, research institutions, Health & Planning and Development Departments and representatives from UN and international and national non-government organizations participated to review the previous DGs, identify gaps and suggest how new DGs should be developed for the country. In the light of participants' suggestions, a framework for PDGN was developed. A consultative workshop was convened to review and refine the framework of the PDGN. Extensive literature search on food consumption patterns, nutrients intake, physical activity, lifestyle, recommended dietary allowances, diet and disease, nutrition and health problems was carried out along with critical evaluation of literature to fully understand the existing situation of dietary intake, the relationship between diet and disease, nutritional and health status of the population. The literature provided necessary inputs and up-to-date snapshot of the nutrition and dietary practices of the population and the difficulties and challenges that could arise during the development of PDGN.

On the basis of existing dietary patterns, nutrient intakes, dietary recommendations, prevalence of malnutrition and diseases in the country, the dietary guidelines were formulated while keeping in view the dietary recommendations and guidelines of WHO/FAO⁹¹ to promote healthy dietary practices and to prevent the rising incidence of overweight and obesity, diabetes, hypertension, cardiovascular diseases, cancers, rickets, arthritis, osteoporosis and other diet related diseases.

A consensus building workshop was conducted to which technical experts, planners and policy makers from universities, research institutions, Ministries of PD&R, Health, Provincial Departments, Azad Jammu and Kashmir and Gilgit-Baltistan were invited. A comprehensive presentation on dietary guidelines was given and working groups were constituted for deliberation and eliciting their inputs. Dietary guidelines were critically reviewed and modified to incorporate the relevant recommendations.

PAKISTAN DIETARY GUIDELINES FOR BETTER NUTRITION (PDGN)

PDGN are a set of dietary recommendations based on scientific and epidemiological evidence to meet the nutrients requirement of population through improved dietary practices and healthy lifestyles. PDGN aims to improve the nutritional status of the population by ensuring that sufficient amounts of nutrients are provided to the body by intelligent and informed food choices and adoption of healthy lifestyles and behavioural practices. Food based interventions are safe risk free, non-invasive, cost effective, sustainable, culturally acceptable and far less expensive as compared to therapeutic treatments. The data suggest that dietary patterns of the Pakistani population in general are inappropriate, unhealthy and inadequate both in terms of quality and quantity to meet the body needs. Nutrients intake data also reveal that diets are unable to provide adequate amount of essential macro and micronutrients to the body as reflected by the higher prevalence of underweight, wasting, and stunting combined with widespread nutritional deficiencies in

women and children¹. These deficiencies fall under the category of public health problems that need to be reversed through integrated, comprehensive dietary interventions (development and dissemination of food-based dietary guidelines, nutrition education, dietary counseling and behavioural change communication).

Health status indicators also highlight that there has been surge in communicable and non-communicable diseases affecting almost all age groups of the population. Women and children are more vulnerable to nutritional and health problems owing to their increased requirements and decreased intakes and less access to health care facilities, poverty and unawareness. Prevalence of overweight and obesity, hepatitis, malaria, tuberculosis, diabetes, hypertension, cardiovascular diseases and cancer is multiplying with the passage of time that tend to have far reaching social, psychological and economic consequences on the society and the country.

Food and food groups

Food is essential for providing nutrients like carbohydrates, proteins, fats, vitamins, minerals and other beneficial substances vital for human survival. None of the foods is capable of providing all the nutrients in an adequate amount to meet the body's requirements. Eating a variety of foods in adequate amount is the only way of nourishing the body. On the basis of nutritional attributes, food may be divided into six groups that include cereals, vegetables, fruits, milk and milk products, meat and pulses, fats and oils.

Each food group has a unique nutrients composition that is different in quality and quantity from other food groups. Fats and oils are included in the food group because they are rich source of energy and essential fatty acids for the body.

Cereals

Cereals include wheat, oats, rice, rye, barley, millet and corn. These grains can be cooked and eaten whole, ground into flour to make a variety of cereal foods like bread, chapatti, nan, pasta and noodles, or made into ready-to-eat breakfast cereals. Wheat, rice and maize are the most commonly consumed cereals in Pakistan. They are the main source of energy, carbohydrates, dietary fibre, protein, vitamins and minerals. Cereals provide over 50% of total energy and protein per day to human body in Pakistan⁵⁴.



Energy balance is important to prevent both under-nutrition and over-nutrition and maintaining healthy body weight. Dietary fibre is essential to maintain body cholesterol level, reduce faecal transit time with

rapid excretion of toxic substances and prevent the incidence of colorectal cancer. Cereals and cereal-based food products are energy dense and thus inadequate or excess intakes are harmful to health.

Vegetables

Vegetables include leafy and non-leafy vegetables with different colours (green, yellow and red) and properties. The leafy vegetables include dark green vegetables (broccoli, spinach, lettuce), red and orange vegetables (tomatoes, carrots, and squash), beans and peas (black beans, split peas, and soybeans). The non-leafy vegetables and tubers include potato, carrots, turnips, reddish, garlic, ginger, onion etc. Vegetables are rich in vitamins, minerals, dietary fibre, water and phytochemicals that protect body from oxidative damage and infectious and chronic diseases. Vegetables are classified as protective foods as they protect body from infections and chronic diseases.



Fruits

Fruits in this group include fresh, frozen, canned and dried fruits. They are a good source of energy, dietary fibre, potassium, vitamins and minerals. They also contain phytochemicals and protect the body from oxidative damage and reduce the risk of cancers and other chronic diseases. Fruits are also classified as protective foods as they protect body from infections and chronic diseases.



Milk and milk products


Milk and milk products include milk, yogurt, cheese, ice cream, custard and other milk based foods. They are good sources of protein, calcium, phosphorus, magnesium, vitamins and minerals.

Adequate amount of milk and milk products are recommended to prevent incidence of rickets, osteomalacia and osteoporosis later in life which occur due to deficiency/low intake of vitamin D, calcium and phosphorus.



Meat and pulses

Meat and pulses include all kinds of meat and organs, fish, eggs and pulses (chick pea, red beans, lentil, mash, mung beans) that are rich in protein and are considered body-building foods. Besides protein, they also provide vitamins and minerals. Adequate amount of meat, fish, eggs and pulses should be taken to reduce and control kwashiorkor, stunting, anaemia and zinc deficiencies.



Fats and oils

Fats and edible oils are an essential part of the Pakistani diet. Fats and oils are major sources of energy, essential for the absorption of fat soluble vitamins and providing essential fatty acids. They are composed of fatty acids and classified into saturated and unsaturated fatty acids. Saturated fatty acids are predominantly present in animal fats (butter and ghee), hydrogenated fat (ghee) and vegetable oils (coconuts and red palm oil). High intake of saturated fatty acids results in synthesis of bad fat i.e., low density lipoprotein (LDL) in the body which has been identified as a risk factor for cardiovascular diseases. Vegetable oils (soybean, sunflower, corn, olive, rapeseed, and sesame seed) are high in unsaturated fatty acids and considered superior over saturated fatty acids as the former promotes the synthesis of good fat i.e., high density lipoprotein (HDL) in the body.



Fats and oils are also divided into essential and non-essential fatty acids. Linolenic and linoleic acids are essential fatty acids not synthesized by the body and must be supplied through the dietary sources as they are vital for performing a variety of physiological functions i.e., cell membrane, development and functioning of the brain and nervous systems. They play an important role in suppressing inflammation and prevent degenerative changes in the body. The composition of fatty acids determines the quality of fats and oils. Fatty acids are also divided into cis and trans fatty acids, the former promotes synthesis of good fat (high density lipoprotein) and the later leads to bad fat (low density lipoprotein) production in the body. Trans fatty acids are produced when oil is transformed into solid fat by industrial processing or when oil and fat are overheated or fatty meat is cooked and eaten. Deep fried foods, bakery products, processed foods and fast foods are potential sources of trans fatty acids which increase low density lipoproteins, modify other lipoproteins and are associated with increased risk of cardiovascular diseases.

Fatty acids are also classified into omega-3, omega-6 and omega-9 on the basis of the location of double bonds in the fatty acid chain. Alpha-linolenic fatty acid is omega-3 essential fatty acid that is converted into docosahexaenoic acid (DHA) and eicosapentaenoic acid (EPA) in the human body that are essential for brain and nervous system functions. Linoleic acid is omega-6 fatty acid found in vegetables oils, grains and nuts while oleic, stearic and erucic fatty acids are omega-9 non-essential fatty acids. The acceptable macronutrient distribution range (AMDR) from fat in the form of total calories is 20-40% for different age groups⁹². A list of foods rich in omega-3, omega-6 and omega-9 fatty acid are given as Annexure-III.

Recommended dietary allowances

Recommended dietary allowances (RDAs) refer to the average daily intake level of a particular nutrient that is likely to meet the nutrient requirements of 97-98% of healthy individuals in a particular life stage or gender group. They are derived from epidemiological studies, which are based on determining mean nutrients intake of the reference population i.e., large nationally representative sample of population to which two standards deviations are added to balance an individual difference and set the nutrients requirement for the general population with an exception to energy requirements which are set equivalent to average mean energy expenditure of a healthy reference population.

Knowledge of nutrients requirement by different age groups is essential for transforming nutrients into portions of foods required by each age group. Nutrients requirement by different age groups are described below to facilitate the understanding of developing PDGN. Recommended dietary allowances for Pakistani population as given in the Food Composition Table for Pakistan⁹³ are reproduced as Annexure-IV.

Infants (0-6 months)

Early infancy covers the period from birth to 6 months, characterized by rapid body growth and development. During this period, infants on an average gain about 4-5 kg weight and 15-16 cm length²¹. Exclusive breastfeeding is recommended due to its unique nutrients composition best suited for the baby. Nutrients supplied by mother's milk are considered adequate for the optimal growth and development of infants. For a particular nutrient, the average intake of full-term infants born to healthy and

well-nourished mothers and are exclusively breastfed has been adopted as the primary basis for deriving the Adequate Intake (AI) for most nutrients during the first 6 months of life. Energy is required for basal metabolism, growth and physical activity which are highest during the first six months starting from 113 kcal/kg/day at birth to 81kcal/kg/day at 6 months for boys and 107 kcal/kg/day at birth to 82 kcal/kg/day at 6 months for girls⁵⁷. Similarly, protein and other nutrients demand per kg body weight are highest during this period to meet the physiological needs and growth of the body.

Infants and young children (6-24 months)

During the second half of infancy and onwards, growth velocity becomes relatively slower with increasing age. Children during this period on an average gain about 4 kg body weight and 20 cm height, therefore require adequate amounts of breast milk and complementary foods to meet the body needs for optimal growth and development²¹. Energy requirements during this age group are estimated between 80-82 kcal/kg/day for girls and boys, respectively⁵⁷. Protein requirements during the first and second years of life have been estimated at 1.69 g, 1.25 g / kg /day, respectively⁹⁴. WHO recommends that complementary foods should be introduced at 6 months of age in addition to breast milk, initially 2-3 times a day for infants from 6-8 months, increasing to 3-4 times daily for infants 9-11 months and 3-4 times complementary foods plus additional nutritious snacks 1-2 times/day for 12-24 month-old children⁹⁵. In terms of energy requirements from complementary feeding during different time periods, it is recommended that complementary foods should provide 200 Kcal/day to 6-8 month-old infants, 300 kcal/day to 9-11month-old infants and 550 kcal/day to 12-23 month-old children⁹⁶. The IYCF recommendations proposed by UNICEF/WHO^{15,18} are given as Annexure-V. For further details, Pakistan Infant and Young Child Strategy, 2016-20 can be consulted⁹⁷.

Children (24-36 months)

Children at this age group grow, learn and acquire skills, enjoy greater mental and motor skills, and are actively engaged in eating and playing. It is a period of learning, talking, communicating, feelings and emotional development. Children by this age begin to explore life independently. It is a time when children learn, develop and display variations in eating behavior, skills, knowledge and attitude relating to food. Energy requirements for children at this age group are about 80 kcal/kg/day⁵⁷ while protein requirements are estimated at 1.1g/kg/day with an acceptable macronutrient distribution range for protein between 5-20% of total energy⁹⁸. Depending on age, size, sex and physical activity total energy requirements of children in this age group range between 1000-1400 kcal/day⁹⁹.



Children (3-10 years)

Growth and development of children 3-10 years continue at a moderate pace but there is a continuous demand of nutrients to meet the body needs. In general, children gain about 2-2.5 kg body weight/year and 5-6cm height/year⁹⁴. Mean body weight and length of boys is generally greater than their corresponding age girls. Energy requirements for boys and girls are estimated at 80 kcal/kg/day and 77 kcal/kg/day at age 3 but decreased to 67 kcal/kg/day and 61 kcal/kg/day at age 10 for boys and girls, respectively⁵⁷. Recommended protein requirements for children 3-10 years are estimated at 1.2 g/kg/day⁹⁴.

Adolescents (10-19 years)

Adolescence period is accompanied by increased physiological and psychological changes with rapid physical growth and development that require increased amounts of nutrients to fulfill the body requirements. At puberty and at peak weight and height accretion, adolescents gain about 3-5 kg/year weight and 6-7 cm/year height, respectively¹⁰⁰. After the age of 17 years, weight and height gain of adolescents reduces substantially to 1.5-2 kg/year and 0.5-1 cm/year, respectively. Adolescents by and large attain 50% of adult weight and 25% of adult height for which optimal nutrition is essential³¹. Energy requirements for adolescent boys and girls are estimated at 55-65 kcal/kg/day and 45-60 kcal/kg/day, respectively. Recommended protein requirements for adolescents are 1.1-1.2 g/kg/day⁹⁴.

Adults (19-60 years)

In adults (age 19 and above), physical growth and development almost comes to an end except skeletal growth that continues with increase in bone mass till the age of 30-35 years. Beyond this period, bone resorption dominates over bone formation process and thus there is a loss of calcium hydroxyapatite from bones and teeth with an increased risk of bone related diseases and fractures. Mean energy expenditure decreases and so does the nutritional requirements. For adults with 60 kg body weight a basal metabolic rate (BMR) of 1515 and sedentary, moderate to heavy or vigorous levels of physical activity, the estimated energy requirements are 2318, 2727 and 3485 kcal/day while for women with 55 kg body weight a BMR of 1241 and sedentary, moderate to heavy or vigorous levels of physical activity, the estimated energy requirements are 1899, 2234 and 2854 kcal/day. Recommended protein requirements for adults are 1 g/kg/day⁹⁴.

Elderly (>60 years)

Nutritional requirements of the elderly vary due to their varying health status. An age related decline in physiological functions combined with the prevalence of chronic diseases and limited health care make the elderly more vulnerable to nutritional deficiencies. More individualized attention is required at this stage to provide the elderly with adequate dietary and nutritional supplements to improve their nutritional well-being.

Pregnancy

Additional nutrients requirements for pregnancy are based on the amount of tissue deposits, fat accumulation and increase in blood volume. Based on the growth and development of the fetus, nutrient requirements vary in three trimesters, minimum additional nutrients requirements during the first trimester, moderate in the second trimester and highest in the third trimester. The total cost of energy during pregnancy is estimated at around 80,000 kcal, which comes to 300 kcal per day¹⁰¹. The RDAs for additional energy and protein intakes during the second and third trimesters are 300 kcal/day, 7 g/day and 23 g/day, respectively¹⁰¹. Optimum weight gain during pregnancy is marker of birth weight; both under and over weight gains during pregnancy are associated with pregnancy complications and adverse pregnancy outcomes. For overweight and obese women- lesser gestational weight gain (<12 kg), for normal- moderate weight gain (12 kg) and for underweight- greater weight gain (12-14 kg) is recommended⁹⁴.

Lactation

Lactation imposes additional nutrients demand on women to meet the requirements of breastfeeding. It has generally been agreed that the nutrients cost incurred on breast feeding shall be met through dietary sources so that nutritional well-being of women and children could be protected to reduce the risk of malnutrition, morbidity and mortality in both groups. Additional energy recommendations for lactating women are 500kcal/day, while 19 g and 13 g additional protein intakes are recommended during the first and second six months of lactations⁹⁴.

MY PLATE

There is no single food that provides all the essential nutrients to the body in adequate amounts for growth and development of infants beyond six months of age. On the basis of quality and characteristics, foods are divided into six groups: (i) Cereals; (ii) Meat and pulses (meats/eggs/pulses); (iii) Dairy (milk and milk products); (iv) Vegetables; (v) Fruits and (vi) Fats and oils. Fats and oils are included in the plate, as they constitute an integral component of Pakistani diet. These groups are presented in My plate form to reinforce that the diet must contain all food groups to satisfy the nutritional requirements of the body.



Cereals group is essential for providing carbohydrates, protein, vitamin B1, vitamin E and minerals. It is the main staple providing over 50% of energy, protein, vitamins and mineral requirements of the body. The acceptable macronutrient distribution range (AMDR) from carbohydrates in the form of total calories is 45-65% for different age groups⁹².

Meat and pulses group includes mutton, beef, poultry, fish, liver, eggs and pulses that are rich sources of proteins and minerals. The AMDR from meat and pulses group in the form of total calories is 5-35% for different age groups⁹².

Dairy group includes milk and milk products essential for skeletal growth and development, neurological and muscular functions. It is a rich source of calcium, phosphorous, protein, lactose, energy and vitamins. Vegetables are essential for providing antioxidants, dietary fibre, vitamins and minerals that protect body

against chronic diseases. Increase consumption of vegetables is recommended to meet the nutritional requirements, maintain desirable body weight, metabolic functions and prevent the incidence of chronic diseases.

Fruits are essential not only for providing vitamins and minerals but also for supplying phyto-chemicals and dietary fibre for optimal metabolic functions, growth and development. Fat, oil, salt and sugar may be used moderately to reduce the risk of overweight and obesity, hypertension, cardiovascular diseases and diabetes.

Portion sizes and frequency of servings

Since nutritional requirement varies by age and gender, emphasis is laid on the development of PDGN by various age groups and gender. For the purpose of PDGN, the population is divided into following age groups:

- Infants (0-6 months)
- Infant and young children (6-24 months)
- Children (24-36 months)
- Children (3-10 years)
- Adolescents (10-19 years)
- Adults (19-60 years)
- Elderly (>60 years)
- Pregnant women
- Lactating mothers

Infants (0-6 months)

It has been well established that breast milk is a complete food for the first six months of life and provides all the essential nutrients in balanced amounts to the baby for meeting its physiological, optimal growth and development requirements. Mothers shall be encouraged, educated and facilitated to adopt exclusive breastfeeding during the first six months.

Infants and young children (6-24 months)

The period between 6-24 months is crucial for laying a strong nutrition base to meet the demands of accelerated growth and development, and to prevent growth faltering, children morbidity and mortality. Exclusive breastfeeding is unable to meet the nutritional requirements of infants beyond six months of age. To complement the baby's nutrients requirements, introduction of soft nutrients dense foods (thick cereals porridge, banana, potato), soups (corn, vegetables, meat) and fresh fruit or fruit juice in small amounts with a gradual increase, along with breastfeeding is vital for child's growth and development. It shall be continued till two years of age after which a gradual shift to the family food is advised.

It is also important that complementary foods must be safe, age-appropriate and consist of a variety of foods to provide adequate amounts of nutrients to the body. The consistency of complementary foods should change from semi-solid to solid as the infants grow. Fortified complementary foods may be given when availability of local diversified foods becomes an issue.

It has been recommended that infants aged 6-8 months should receive complementary foods 2-3 times/day, infants aged 9-11 months should receive complementary foods 3-4 times/day and additional nutritious snacks 1-2 times/day for children older than 1 year⁹⁵. The estimated energy requirements of complementary foods for infants between 6-8 months are 200 kcal/day; 300 kcal/day for 9-11 months and 550 kcal/day for 12-24 month-old children⁹⁶.

Dietary recommendations for infants and young children (6-24 months)

- *Exclusive breast feeding from 0 to 6 months.*
- *Continue breast feeding along with complementary foods adequate in quality and quantity (2-4 times/day) from a variety of soft foods starting from liquid to semi-solid and solid foods gradually suitable for young children under 2 years of age.*
- *Salt and sugar should not be added to complementary foods¹⁰².*
- *Provide nutrients dense snacks from fresh foods (1-2 times/day) as per child's appetite and requirement.*

Some of the commonly consumed complementary foods given in the Food Composition Table for Pakistan⁹³ are reproduced as Annexure-VI.

Children (24-36 months)

Children in this age group require a well-balanced, nutrient dense diet to meet the nutritional requirements of the body needed for the maintenance of basal metabolic rate, optimal physical and mental growth and development, and increased levels of physical activity. The parents and caregiver must help children in establishing healthy dietary habits, good hygiene practices, and healthy behavior and lifestyle attitudes for an improved and productive life. The basic principle of meal planning should be that diet must be inclusive of the basic five food groups and quantitatively and qualitatively sufficient to meet the growth and development demands of the body.

Food and portion sizes for children (24-36 months)

| Food Group | No. of Servings /day | Portion size and description |
|------------------------|----------------------|---|
| Milk and milk products | 2 | 1 serving=1cup of milk or 1 cup of yogurt or 1 slice of cheese or 1 cup of kheer or feerni or other milk-based products equivalent to nutrients supplied by 1 cup of milk. 1 cup of whole milk will provide 15g carbohydrates, 6 g protein, 8 g fat and 150 calories. |
| Cereals | 1-2 | 1 serving= 2 slice of bread (toast) or 1 chapatti or 1 cup of cooked rice or 1 cup of cereals equivalent to nutrients supplied by 2 slice of bread. One serving of cereals bread=(2 toast x 28 g=56 g) shall provide 30 g carbohydrates, 6 g protein, 0-2 g fat and 160 Kcal. |
| Vegetables | 1-2 | 1 serving= ½ cup of cooked non-starchy vegetables or ½ cup of vegetables juice/soup or 1 cup of fresh vegetables/salad. One serving of vegetables will provide 5 g carbohydrates, 2 g protein and 25 calories. One serving of starchy vegetables (1 potato (100 g) or maize (1/2 cup) or peas green ½ cup) will provide 15 g carbohydrates, 3 g protein, 0-1 g fat and 80 calories. |
| Fruits | 1-2 | 1 serving=1 medium size banana or 1 apple or 1 peach or 2-3 plums or 3-4 apricots. Each serving will provide 20 g carbohydrates and 80 calories. |
| Meat and pulses | 1-2 | 1 serving of meat (28 g lean meat=2-3 small pieces of meat or 1-2 pieces of fish or 1 egg, or ½ cup of cooked pulses). One serving of lean meat will provide 7 g protein, 3 g fat and 55 calories. One serving of meat with medium fat will provide 7 g protein, 5 g fat and 75 calories. One serving of meat with high fat will provide 7 g protein, 8 g fat and 100 calories. |

Salt, Sugar and Oil

- Use minimum amount of salt, sugars and oils in cooking use less than 5 g salt/day, less than 10% calories from free sugars and less than 10% of total fat calories from saturated fat.
- Avoid junk and refined foods, sweets, soft drinks, french fries, samosa and other deep fried foods.
- Use clean water for washing vegetables and fruits and protect foods from flies, pathogens, dirt, and cover and refrigerate them at 5°C or below for a day only.

Children (3-10 years)

Growth and development in children between 3-10 years of age, gets slower than infancy and is considered a period of catch-up growth to cover the nutritional deficit gaps and build up the deficient body nutritional stores. It is an opportune time to improve nutritional status and to boost body's immune system for fighting against infections and diseases. Children at this age develop better taste acuity, improved digestive and absorptive capacities, enhanced enzymes and hormonal secretion, coordination and control to digest, absorb and utilize family foods. This period is also important for learning and acquiring skills and developing habits for eating (adult type) family foods. At this stage a child picks and imitates what their peers and older people eat and drink. Encouraging children to adopt healthy eating practices pay dividends to rest of their life.

Introducing a variety of nutrients dense foods that comprise of milk and milk products, meat and meat products, vegetables and fruits, cereals and pulses with legumes and increasing meal frequencies will enable children not only to meet their daily nutritional requirements but also to replete the nutrient stores of the body.

Food and portion sizes for children (3-10 years)

| Food Group | No. of Servings /day | Portion size and description |
|------------------------|----------------------|---|
| Milk and milk products | 2-3 | 1 serving=1cup of milk or 1 cup of yogurt or 1 slice of cheese or 1 cup of kheer or feerni or other milk-based products equivalent to nutrients supplied by 1 cup of milk. 1 cup of whole milk will provide 15 g carbohydrates, 6 g protein, 8 g fat and 150 calories. |
| Cereals | 2-4 | 1 serving= 2 slice of bread (toast) or 1 chapatti or 1 cup of cooked rice or 1 cup of cereals equivalent to nutrients supplied by 2 slice of bread. One serving of cereals bread=(2 toast x 28 g=56 g) shall provide 30 g carbohydrates, 6 g protein, 0-2 g fat and 160 Kcal. |
| Vegetables | 1-2 | 1 serving= ½ cup of cooked non-starchy vegetables or ½ cup of vegetables juice/soup or 1 cup of fresh vegetables/salad. One serving of vegetables will provide 5 g carbohydrates, 2 g protein and 25 calories. One serving of starchy vegetables (1 potato (100 g) or maize (1/2 cup) or peas green ½ cup) will provide 15 g carbohydrates, 3 g protein, 0-1 g fat and 80 calories. |
| Fruits | 1-2 | 1 serving=1 medium size banana or 1 apple or 1 peach or 2-3 plums or 3-4 apricots. Each serving will provide 20 g carbohydrates and 80 calories. |
| Meat and pulses | 1-2 | 1 serving of meat (28 g lean meat=2-3 small pieces of meat or 1-2 pieces of fish or 1 egg, or ½ cup of cooked pulses). One serving of lean meat will provide 7 g protein, 3 g fat and 55 calories. One serving of meat with medium fat will provide 7 g protein, 5 g fat and 75 calories. One serving of meat with high fat will provide 7 g protein, 8 g fat and 100 calories. |

Salt, Sugar and Oil

- Use minimum amount of salt, sugars and oils in cooking use less than 5 g salt/day, less than

10% calories from free sugars and less than 10% of total fat calories from saturated fat.

- Avoid junk and refined foods, sweets, soft drinks, french fries, samosa and other deep fried foods.
- Use clean water for washing vegetables and fruits and protect foods from flies, pathogens, dirt, and cover and refrigerate them at 5°C or below for a day only.

Adolescents (10-19 years)

Adolescence is a period of rapid growth and development marked by rapid changes in body size, shape, and composition that require adequate nutrition for attaining optimum weight, height, replenish nutrient stores to reach full genetic potential in physical, mental and scholastic performance. The rapid growth is accompanied by physiological changes that include increased hormonal secretions and reproductive development with secondary characteristics, cell proliferation and increased blood volume, which increases the nutrient requirements. At the peak of the adolescent growth spurt, the nutritional requirements may be twice as high as those of the remaining period of adolescence¹⁰³.

During adolescence, weight and height increases by about 50% and 20% of adult weight and height respectively, that puts greater demand of nutrients which could be only met through consumption of nutrients dense diversified foods³¹. However, girls' marriage during adolescence followed by subsequent pregnancies and lactation results in additional nutritional stress on women making them vulnerable to nutritional deficiencies with increased risk of morbidity, neonatal and maternal mortalities.

Good and adequate nutrition during adolescence are pre-requisite for health, productivity and economic outputs. Adolescent girls need special attention with regard to nutritional well-being because of their multi-dimensional responsibilities at home and outdoors including bringing water, collecting fire wood, and helping mothers in home in domestic chores and rearing animals. Failing to cope with increasing nutrient's demand, many adolescents remain underweight and stunted which has been attributed to obstetric complications and adverse pregnancy outcomes¹⁰⁴.

The vicious cycle of malnutrition perpetuates from generation to generation with increased prevalence of diseases, overburden on health systems and substantial social and economic losses for the country. Nutrition during all stages of life but particularly during adolescence period is crucial and cost effective as the amount spent on foods is negligible to costs incurred on treatments and sufferings caused by diseases, disabilities, loss of work and premature deaths in later life.

Food and portion sizes for adolescents (10-19 years)

| Food Group | No. of Servings /day | Portion size and description |
|------------------------|----------------------|--|
| Milk and milk products | 3-4 | 1 serving=1cup of milk or 1 cup of yogurt or 1 slice of cheese or 1 cup of kheer or feerni or other milk-based products equivalent to nutrients supplied by 1 cup of milk. 1 cup of whole milk will provide 15 g carbohydrates, 6 g protein, 8 g fat and 150 calories. |
| Cereals | 5-6 | 1 serving= 2 slice of bread (toast) or 1 chapatti or 1 cup of cooked rice or 1 cup of cereals equivalent to nutrients supplied by 2 slice of bread. One serving of cereal bread=(2 toast x 28 g=56 g) shall provide 30 g carbohydrates, 6 g protein, 0-2 g fat and 160 Kcal. |
| Vegetables | 2-3 | 1 serving= ½ cup of cooked non-starchy vegetables or ½ cup of vegetables juice/soup or 1 cup of fresh vegetables/salad. One serving of vegetables will provide 5 g carbohydrates, 2 g protein and 25 calories. One serving of starchy vegetables (1 potato (100 g) or maize (1/2 cup) or peas green ½cup) will provide 15 g carbohydrates, 3 g protein, 0-1 g fat and 80 calories. |
| Fruits | 2-3 | 1 serving=1 medium size banana or 1 apple or 1 peach or 2-3 plums or 3-4 apricots. Each serving will provide 20 g carbohydrates and 80 calories. |
| Meat and pulses | 2-3 | 1 serving of meat (28 g lean meat=2-3 small pieces of meat or 1-2 pieces of fish or 1 egg, or ½ cup of cooked pulses). One serving of lean meat will provide 7 g protein, 3 g fat and 55 calories. One serving of meat with medium fat will provide 7 g protein, 5 g fat and 75 calories. One serving of meat with high fat will provide 7 g protein, 8 g fat and 100 calories. |

Salt, Sugar and Oil

- Use minimum amount of salt, sugars and oils in cooking, use less than 5 g salt/day, less than 10% calories from free sugars and less than 10% of total fat calories from saturated fat.
- Avoid junk, refined foods, sweets, soft drinks, french fries, samosa and other deep fried foods.
- Use clean water for washing vegetables and fruits and protect foods from flies, pathogens, dirt, cover and refrigerate them at 5 °C or below for a day only.

Adult (19-60 years)

Adulthood is a period where growth and development to a large extent cease, however, skeletal development and deposition of calcium, phosphorus and magnesium in bones continues till the age of 30-35 years. Beyond this age, bone resorption predominates over bone formation and the time taken to deplete skeletal mineral deposits is proportional to the amount of mineral deposits, the higher are the mineral deposits during skeletal growth and development, the greater would be the time required to erode mineral deposits from bones.

Good and adequate nutrition in combination with regular aerobic exercise and physical activities are essential throughout adulthood period to maintain optimum nutritional status, prevent nutritional deficiencies, overweight and obesity which are the leading causes of non-communicable diseases including diabetes, hypertension, cardiovascular diseases and cancers. In recent years, prevalence of overweight and obesity in adults has been increasing in an unprecedented manner resulting in increased morbidity and mortality. Dietary modifications including reduction of sugars, salt, saturated fats, trans-fatty acids, increased physical activity and healthy lifestyles are cost effective interventions essential to reduce and control adulthood morbidities and mortalities.

Food and portion sizes for adults (19-60 years)

| Food Group | No. of Servings /day | Portion size and description |
|------------------------|----------------------|---|
| Milk and milk products | 2-3 | 1 serving=1cup of milk or 1 cup of yogurt or 1 slice of cheese or 1 cup of kheer or feerni or other milk-based products equivalent to nutrients supplied by 1 cup of milk). 1 cup of whole milk will provide 15 g carbohydrates, 6 g protein, 8 g fat and 150 calories. |
| Cereals | 4-5 | 1 serving= 2 slice of bread (toast) or 1 chapatti or 1 cup of cooked rice or 1 cup of cereals equivalent to nutrients supplied by 2 slice of bread. One serving of cereal bread=(2 toast x 28 g=56 g) or equivalent amount of other cereals shall provide 30 g carbohydrates, 6 g protein, 0-2 g fat and 160 Kcal. |
| Vegetables | 2-3 | 1 serving= ½ cup of cooked non-starchy vegetables or ½ cup of vegetables juice/soup or 1 cup of fresh vegetables/salad). One serving of vegetables will provide 5 g carbohydrates, 2 g protein and 25 calories. One serving of starchy vegetables (1 potato (100 g) or maize (1/2 cup) or peas green ½cup) will provide 15 g carbohydrates, 3 g protein, 0-1 g fat and 80 calories. |
| Fruits | 2-3 | 1 serving=1 medium size banana or 1 apple or 1 peach or 2-3 plums or 3-4 apricots. Each serving will provide 20 g carbohydrates and 80 calories. |
| Meat and pulses | 2-3 | 1 serving of meat (28 g lean meat=2-3 small pieces of meat or 1-2 pieces of fish or 1 egg, or ½ cup of cooked pulses). One serving of lean meat will provide 7 g protein, 3 g fat and 55 calories. One serving of meat with medium fat will provide 7 g protein, 5 g fat and 75 calories. One serving of meat with high fat will provide 7 g protein, 8 g fat and 100 calories. |

Salt, Sugar and Oil

- Use minimum amount of salt, sugars and oils in cooking, use less than 5 g salt/day, less than 10% calories from free sugars and less than 10% of total fat calories from saturated fat.
- Avoid junk, refined foods, sweets, soft drinks, french fries, samosa and other deep fried foods.
- Use clean water for washing vegetables and fruits and protect foods from flies, pathogens, dirt, cover and refrigerate them at 5°C or below for a day only.

Elderly (>60 years)

Older people have special dietary and health care needs to meet the body nutrients requirements, boost their immune systems and maintain optimum level of physical activity to prevent the advancement of diseases and complications that may increase the risk of falling, bone fractures and immobility. Diet adequate both in quality and quantity is vital for nutrition wellbeing and prevent the incidence and severity of diseases.

Elderly need to enhance dietary and health care to ward off non-communicable and degenerative diseases owing to age and compromised nutrition and health status. Impaired taste acuity, gastrointestinal, enzymatic, hormonal and metabolic functions in older persons make them more vulnerable to malnutrition and diseases. Qualitative and quantitative dietary modification is essential to provide adequate amount of dietary fibre, proteins, vitamins, minerals, water and fluid to prevent constipation and other age related diseases.

Food and portion sizes for elderly (>60 years)

| Food Group | No. of Servings /day | Portion size and description |
|------------------------|----------------------|---|
| Milk and milk products | 2-3 | 1 serving=1cup of milk or 1 cup of yogurt or 1 slice of cheese or 1 cup of kheer or feerni or other milk-based products equivalent to nutrients supplied by 1 cup of milk. 1 cup of whole milk will provide 15 g carbohydrates, 6 g protein, 8 g fat and 150 calories. |
| Cereals | 4-5 | 1 serving= 2 slice of bread (toast) or 1 chapatti or 1 cup of cooked rice or 1 cup of cereals equivalent to nutrients supplied by 2 slice of bread. One serving of cereal bread =(2 toast x 28 g=56 g) shall provide 30 g carbohydrates, 6 g protein, 0-2 g fat and 160 Kcal. |
| Vegetables | 2-3 | 1 serving= ½ cup of cooked non-starchy vegetables or ½ cup of vegetables juice/soup or 1 cup of fresh vegetables/salad). One serving of vegetables will provide 5 g carbohydrates, 2 g protein and 25 calories. One serving of starchy vegetables (1 potato (100 g) or maize (1/2 cup) or peas green ½cup) will provide 15 g carbohydrates, 3 g protein, 0-1 g fat and 80 calories. |
| Fruits | 2-3 | 1 serving=1 medium size banana or 1 apple or 1 peach or 2-3 plums or 3-4 apricots. Each serving will provide 20 g carbohydrates and 80 calories. |
| Meat and pulses | 2-3 | 1 serving of meat (28 g lean meat=2-3 small pieces of meat or 1-2 pieces of fish or 1 egg, or ½ cup of cooked pulses). One serving of lean meat will provide 7 g protein, 3 g fat and 55 calories. One serving of meat with medium fat will provide 7 g protein, 5 g fat and 75 calories. One serving of meat with high fat will provide 7 g protein, 8g fat and 100 calories. |

Salt, Sugar and Oil

- Use minimum amount of salt, sugars and oils in cooking, use less than 5 g salt/day, less than

10% calories from free sugars and less than 10% of total fat calories from saturated fat.

- Avoid junk, refined foods, sweets, soft drinks, french fries, samosa and other deep fried foods.
- Use clean water for washing vegetables and fruits and protect foods from flies, pathogens, dirt, cover and refrigerate them at 5 °C or below for a day only.

Pregnant women

Good nutrition during pregnancy is not only essential for maintaining maternal nutritional and health status but also for growth and development of fetus. Adequate and better quality diet during pregnancy is vital for meeting the physiological requirements of expanding maternal tissues and blood volume as well as growth and development of fetus to avoid unfavorable pregnancy outcomes.

Poor nutritional status during pregnancy leads to poor intrauterine growth and development with increased risk of small for gestational age babies, low birth weight babies with impaired physiological functions, increased morbidity and neonatal mortality.

Improving maternal nutrition through improved dietary diversification and nutritional supplements appears to be one of the most profitable investments in paying long term dividends by reducing health problems, medical costs and increasing productivity and economic outputs. Nutritional requirements of pregnant women depend on the pre-pregnancy nutritional status, the poorer the nutritional status prior to conception the higher are the recommended nutrient intakes and weight gain during pregnancy. Pre-pregnancy weight and weight gain during pregnancy has been associated with birth weight of the baby. Both low and high weight gains result in adverse pregnancy outcomes⁹⁴. Since first trimester is limited mostly to embryonic and cell differentiation with nominal weight gain (1-2 kg), additional calorie intake is recommended from second trimester onwards to avoid undesirable maternal weight gain and pregnancy complications. Twelve to sixteen kilograms weight gain is recommended for women with normal body mass index at conception and higher weight gain for women with lower body mass index and lower weight gain for overweight and obese women is recommended. Due to increased growth and development, additional 300 kilocalories, 10 g protein, 15 mg iron, 220 µg folate, 400 mg calcium, 400 mg phosphorus per day is recommended for pregnant women¹⁰¹.

Food and portion sizes for pregnant women

| Food Group | No. of Servings /day | Portion size and description |
|------------------------|----------------------|---|
| Milk and milk products | 2-3 | 1 serving=1cup of milk or 1 cup of yogurt or 1 slice of cheese or 1 cup of kheer or feerni or other milk-based products equivalent to nutrients supplied by 1 cup of milk). 1 cup of whole milk will provide 15 g carbohydrates, 6 g protein, 8 g fat and 150 calories. |
| Cereals | 5-6 | 1 serving= 2 slice of bread (toast) or 1 chapatti or 1 cup of cooked rice or 1 cup of cereals equivalent to nutrients supplied by 2 slice of bread). One serving of cereal bread=(2 toast x 28 g=56 g) shall provide 30 g carbohydrates, 6 g protein, 0-2 g fat and 160 Kcal. |
| Vegetables | 2-3 | 1 serving= ½ cup of cooked non-starchy vegetables or ½ cup of vegetables juice/soup or 1 cup of fresh vegetables/salad). One serving of vegetables will provide 5 g carbohydrates, 2 g protein and 25 calories. One serving of starchy vegetables (1 potato (100 g) or maize (1/2 cup) or peas green ½cup) will provide 15 g carbohydrates, 3 g protein, 0-1 g fat and 80 calories. |
| Fruits | 2-3 | 1 serving=1 medium size banana or 1 apple or 1 peach or 2-3 plums or 3-4 apricots. Each serving will provide 20 g carbohydrates and 80 calories. |
| Meat and pulses | 2-3 | 1 serving of meat (28 g lean meat=2-3 small pieces of meat or 1-2 pieces of fish or 1 egg, or ½ cup of cooked pulses). One serving of lean meat will provide 7 g protein, 3 g fat and 55 calories. One serving of meat with medium fat will provide 7 g protein, 5 g fat and 75 calories. One serving of meat with high fat will provide 7 g protein, 8 g fat and 100 calories. |

Salt, Sugar and Oil

- Use minimum amount of salt, sugars and oils in cooking, use less than 5 g salt/day, less than 10% calories from free sugars and less than 10% of total fat calories from saturated fat.
- Avoid junk, refined foods, sweets, soft drinks, french fries, samosa and other deep fried foods.
- Use clean water for washing vegetables and fruits and protect foods from flies, pathogens, dirt, cover and refrigerate them at 5 °C or below for a day only.

Lactating women

Lactation is a period where nutritional requirements of women are increased due to breast feeding of the new born baby. It has been estimated that mothers during the first six months secrete 750 ml milk/day, which drops to 600 ml milk/day in the second six months of the postnatal period providing about 525 kcal/day and 420kcal/day, respectively⁴⁸. To compensate the energy and micronutrients transferred from the mother to infant through breast feeding and also to enhance the quality of milk and supply, additional 500 kcal, 12-15 g protein, along with optimal amount of micronutrients (vitamin A, vitamin C, vitamin B1, vitamin B2, niacin, vitamin B6, vitamin B12, calcium, vitamin D, phosphorus, zinc, iodine and

selenium) per day are recommended⁴⁸. The lactating women need special attention also to rest well in addition consuming healthy diet, which can help in maintaining good supply of milk as well.

Food and portion sizes for lactating women

| Food Group | No. of Servings /day | Portion size and description |
|------------------------|----------------------|--|
| Milk and milk products | 3-4 | 1 serving=1cup of milk or 1 cup of yogurt or 1 slice of cheese or 1 cup of kheer or feerni or other milk-based products equivalent to nutrients supplied by 1 cup of milk). 1 cup of whole milk will provide 15 g carbohydrates, 6 g protein, 8 g fat and 150 calories. |
| Cereals | 6-7 | 1 serving= 2 slice of bread (toast) or 1 chapatti or 1 cup of cooked rice or 1 cup of cereals equivalent to nutrients supplied by 2 slice of bread. One serving of cereal bread =(2 toast x 28 g=56 g) shall provide 30 g carbohydrates, 6 g protein, 0-2 g fat and 160 Kcal. |
| Vegetables | 2-3 | 1 serving= ½ cup of cooked non-starchy vegetables or ½ cup of vegetables juice/soup or 1 cup of fresh vegetables/salad). One serving of vegetables will provide 5 g carbohydrates, 2 g protein, 0 g fat and 25 calories. One serving of starchy vegetables (1 potato (100 g) or maize (1/2 cup) or peas green ½cup) will provide 15 g carbohydrates, 3 g protein, 0-1 g fat and 80 calories. |
| Fruits | 2-3 | 1 serving=1 medium size banana or 1 apple or 1 peach or 2-3 plums or 3-4 apricots. Each serving will provide 20 g carbohydrates and 80 calories. |
| Meat and pulses | 2-3 | 1 serving of meat (28 g lean meat=2-3 small pieces of meat or 1-2 pieces of fish or 1 egg, or ½ cup of cooked pulses). One serving of lean meat will provide 7 g protein, 3 g fat and 55 calories. One serving of meat with medium fat will provide 7 g protein, 5 g fat and 75 calories. One serving of meat with high fat will provide 7 g protein, 8 g fat and 100 calories. |

Salt, Sugar and Oil

- Use minimum amount of salt, sugars and oils in cooking use less than 5 g salt/day, less than 10% calories from free sugars and less than 10% of total fat calories from saturated fat.
- Avoid junk, refined foods, sweets, soft drinks, french fries, samosa and other deep fried foods.
- Use clean water for washing vegetables and fruits and protect foods from flies, pathogens, dirt, cover and refrigerate them at 5 °C or below for a day only.

An inclusive table indicating number of servings of five food groups for different age and physiological groups is attached as Annexure VII.

SAMPLE MENUS BY AGE AND PHYSIOLOGICAL STATUS

Sample menu for children (3-10 years)

| Time | Food | Portion Size | Calories (Kcal) | Protein (g) |
|-----------|--------------------------------------|---|-----------------|-------------|
| Breakfast | Chapatti/paratha/toast/cereals* | 1-2 toast or ½-1 chapatti or ¼-½ paratha or ½-1 cup cereals | 80-160 | 3-6 |
| | Egg fried‡ | 1 medium | 90 | 6 |
| | Milk† | ½ cup | 75 | 3 |
| Snack | Banana or any other seasonal fruit ¶ | One medium | 80 | ----- |
| | Shami kabab‡ | 1 | 90 | 6 |
| Lunch | Dal§ | ½-1 plate | 70-140 | 4-8 |
| | Roti* | ½-1 chapatti | 80-160 | 3-6 |
| | Yogurt† | ½ cup | 75 | 3 |
| | Salad§ | ½-1 bowl | 13-25 | ----- |
| Snack | Apple or any other seasonal fruit ¶ | 1 | 80 | ----- |
| | Peanuts** | 1 serving | 160 | 7 |
| Dinner | Chicken curry† | 1 plate | 200 | 6 |
| | Roti | ½- 1 chapatti | 80-160 | 3-6 |
| | Salad§ | ½-1 bowl | 13-25 | ----- |
| | Apple or any other seasonal fruit¶ | 1 | 80 | ----- |
| | Milk/Kheer† | ½ cup | 75 | 3 |
| Total | | | 1341-1675 Kcal | 47-60 g |

*Chapatti or paratha or toast or fortified cereals providing a similar amount of calories and proteins;

‡ Egg fried or meat, chicken, fish or lentil or chickpea or red beans or shami kabab or any other meat based protein and pulses providing a similar amount of calories and proteins;

¶Banana or apple or any other seasonal fruit providing a similar amount of calories;

§Chicken curry or dal or meat curry or beans providing a similar amount of calories and proteins;

† ½ cup kheer or ½ cup milk or ½ cup yogurt or ½ ice cream or ½ slice cheese or any other milk-based product providing a similar amount of calories and proteins;

** Peanuts or any other nuts providing a similar amount of calories and proteins



Note: 1 Cup = 250 ml; 1 plate pulao or dal or vegetables (100 g cooked food); 1 bowl fresh salad = (100 g);

1 serving peanuts = 1 ounce peanuts

Sample menu for adolescents (10-19 years)

| Time | Food | Portion Size | Calories (Kcal) | Protein (g) |
|------------------|-------------------------------------|--------------|-----------------|-------------|
| Breakfast | Milk† | 1 cup | 150 | 6 |
| | Egg fried‡ | 1 medium | 90 | 6 |
| | Roti/toast/cereals* | ½-1 chapatti | 80-160 | 8 |
| Snack | Apple or any other seasonal fruit¶ | 1 medium | 80 | ----- |
| | Potato cutlet§ | 1-2 | 49-98 | 1-2 |
| Lunch | Lentil‡ | ½ plate | 70 | 4 |
| | Roti | 1-2 chapatti | 160-320 | 6-12 |
| | Yogurt† | ½ cup | 75 | 3 |
| | Salad§ | ½-1 bowl | 13-25 | ----- |
| Snack | Banana¶ or any other seasonal fruit | 1 medium | 80 | ----- |
| | Peanuts** | 1 serving | 160 | 7 |
| Dinner | Aloo ghost‡ | 1 plate | 174 | 6-8 |
| | Roti | 1-2 chapatti | 160-320 | 6-12 |
| | Salad§ | ½-1 bowl | 13-25 | ----- |
| | Raita† | ½-1 cup | 75-150 | 3-6 |
| | Guava or any other seasonal fruit¶ | 1 medium | 80 | ----- |
| Snack | Carrot halwa† | 1 cup | 250 | 6 |
| Total | | | 1758-2307 Kcal | 57-78 g |

† 1 cup milk or 1 cup kheer or 1 cup yogurt or 1 cup ice cream or 1 slice cheese or any other milk-based product providing a similar amount of calories and proteins;

‡ Egg fried or meat, chicken, fish or lentil or chickpea or red beans or shami kabab or any other meat based curry providing a similar amount of calories and proteins;

*1-2 toast or ½-1 chapatti or ½-1 cup cereals or ¼-½ paratha or any other cereal providing a similar amount of calories and proteins;

¶ Banana or any other seasonal fruit providing a similar amount of calories; §Potato cutlet or starchy vegetables providing a similar amount of calories and proteins;

‡Aloo ghost or chicken curry or any other curry or pulses providing a similar amount of calories and proteins;

†Carrot-halwa or pumpkin halwa or any other milk-based dessert providing a similar amount of calories and proteins;

** Peanuts or any other nuts providing a similar amount of calories and proteins

Note:1 Cup =250 ml; 1 plate pulao or dal or vegetables (100 g cooked food); 1 bowl fresh salad= (100 g);

1 serving peanuts=1 ounce peanuts

Sample menu for adults (19-60 years)

| Time | Food | Portion Size | Calories (Kcal) | Protein (g) |
|------------------|------------------------------------|--------------|-----------------|-------------|
| Breakfast | Roti/toast/cereals* | 1-2 chapatti | 160-320 | 6-12 |
| | Egg fried‡ | 1 medium | 90 | 6 |
| | Apple or any other seasonal fruit¶ | 1 medium | 80 | ----- |
| | Milk† | 1 cup | 150 | 6 |
| Snack | Shami sandwich‡‡ | 1 | 180 | 8 |
| Lunch | Spinach and potato curry§ | 1 plate | 109 | 3 |
| | Roti | 2 chapatti | 320 | 6-12 |
| | Yogurt† | ½ cup | 75 | 3 |
| | Salad§ | 1 bowl | 25 | ----- |
| Snack | Banana or any other seasonal fruit | 1 medium | 80 | ----- |
| Dinner | Pulao** | 1 plate | 250 | 8 |
| | Fish ‡ | 1-2 | 127-254 | 10-20 |
| | Salad§ | ½-1 bowl | 13-25 | ----- |
| | Raita† | ½-1 cup | 75-150 | 3-6 |
| | Apple ¶ | 1 medium | 80 | ----- |
| Snack | Peanuts*** | 1 serving | 160 | 7 |
| | Kheer† | 1 cup | 150 | 6 |
| Total | | | 2124-2498 Kcal | 72-97 g |

*1 roti or 2 toast or ½ paratha or 1 cup fortified cereals providing a similar amount of calories and proteins;

‡ Egg fried or meat, chicken, fish or lentil or chickpea or red beans or shami kabab;

† 1 cup milk or 1 cup kheer or 1 cup yogurt or 1 cup ice cream or 1 slice cheese or any other milk-based product providing a similar amount of calories and proteins;

¶ Banana or apple or any other seasonal fruit providing a similar amount of calories;

‡‡ Shami sandwich or egg sandwich, or chicken sandwich or any other meat or pulses based food providing the similar amount of calories and proteins;

§ Spinach and potato curry or any other vegetable curry providing a similar amount of calories and proteins

** Pulao or dal-roti or vegetable roti providing a similar amount of calories and proteins;

*** Peanuts or any other nuts providing a similar amount of calories and proteins



Note: 1 Cup = 250 ml; 1 plate pulao or dal or vegetables (100 g cooked food); 1 bowl fresh salad = (100 g);

1 serving peanuts = 1 ounce peanuts

Sample menu for elderly (> 60)

| Time | Food | Portion size | Calories (Kcal) | Protein (g) |
|-----------|---|--------------------------|-----------------|-------------|
| Breakfast | Cereals/chapatti/toast* | 1 cup/1 chapatti/2 toast | 160 | 6 |
| | Milk† | 1 cup | 150 | 6 |
| | Egg fried‡ | 1 medium | 90 | 6 |
| | Banana or any seasonal fruit or juice¶ | 1 medium/1 cup | 80 | ----- |
| Snack | Egg sandwich/ chicken sandwich/ or shami sandwich‡‡ | 1 | 180 | 8 |
| Lunch | Spinach curry§ | 1 plate | 109 | 3 |
| | Roti | 2 chapatti | 320 | 12 |
| | Salad± | 1 bowl | 25 | ----- |
| | Yogurt† | 1 cup | 150 | 6 |
| Snack | Shami kabab# | 1 | 90 | 6 |
| | Banana or any other seasonal fruit¶¶ | 1 medium/1 cup | 80 | ----- |
| Dinner | Vegetable rice*** | 1 plate | 187 | 4 |
| | Dal** | 1 plate | 140 | 8 |
| | Salad± | 1 /2 bowl | 13 | ---- |
| | Raita/yogurt† | 1 cup | 150 | 6 |
| | Milk/Kheer/dessert† | 1 cup | 150 | 6 |
| Total | | | 2074 Kcal | 77 g |

*1 cup fortified cereals or 1 roti or 2 toast or ½ paratha providing a similar amount of calories and proteins;
†1 cup milk or 1 cup yogurt or 1 cup kheer or 1 ice cream or any other milk-based product providing a similar amount of calories and proteins;

‡ Egg fried or matar keema or meat, chicken, fish or lentil or chickpea or red beans or any other pulses providing a similar amount of calories and proteins;

¶ Banana or any other seasonal fruit providing a similar amount of calories;

‡‡Egg sandwich or chicken sandwich or shami sandwich or boiled chickpea or red beans or slice of pizza providing a similar amount of calories and proteins;

§Spinach curry or any other vegetable providing a similar amount of calories and protein;

Shami kabab or red beans or boiled gram or meat based snack providing a similar amount of calories and proteins;

** Dal or meat curry providing a similar amount of calories and protein;

± Salad or any other leafy vegetables providing a similar amount of calories;

*** Vegetable rice or bread with salad providing a similar amount of calories and protein;



© FAO/Nomeena Anis

Note:1 Cup =250 ml; 1 plate pulao or dal or vegetables (100 g cooked food); 1 bowl fresh salad= (100 g)

Sample menu for pregnant women

| Time | Food | Portion Size | Calories (Kcal) | Protein (g) |
|-----------|-----------------------------|--------------|-----------------|-------------|
| Breakfast | Roti/toast/cereals* | 1 chapatti | 160 | 6 |
| | Egg fried‡ | 1 medium | 90 | 6 |
| | Milk† | 1 cup | 150 | 6 |
| Snack | Beans or gram boiled‡‡ | 1 cup | 220 | 10 |
| | Guava¶ | 1 medium | 80 | ----- |
| Lunch | Meat ball or kofta curry‡‡‡ | 1 plate | 268 | 8 |
| | Roti* | 2 chapatti | 320 | 12 |
| | Yogurt† | ½ cup | 75 | 3 |
| | Salad§ | 1 bowl | 25 | ----- |
| Snack | Shami sandwich‡‡ | 1-2 | 180-360 | 8-16 |
| Dinner | Vegetable rice** | 1 plate | 187 | 4 |
| | Dal chana‡ | 1 plate | 140 | 8 |
| | Salad§ | 1 bowl | 25 | ----- |
| | Apple ¶ | 1 medium | 80 | ----- |
| Snack | Peanuts*** | 1 serving | 160 | 7 |
| | Carrot halwa**** | 1 cup | 250 | 6 |
| | Milk† | 1 cup | 150 | 6 |
| Total | | | 2560-2740 Kcal | 90-98 g |

*1 roti or 2 toast or ½ paratha or 1 cup fortified cereals providing a similar amount of calories and proteins;

‡ Egg fried or matar keema or meat, chicken, fish or lentil or chickpea or red beans or any other pulses providing a similar amount of calories and proteins;

† 1 cup milk or 1 cup yogurt or 1 cup kheer or 1 ice cream or any other milk-based product providing a similar amount of calories and proteins;

¶ Any seasonal fruit providing a similar amount of calories;

‡‡ Chicken sandwich or shami sandwich or egg sandwich or boiled chickpea or red beans or slice of pizza or peanut butter sandwich providing a similar amount of calories and proteins;

‡‡‡ Meat or kofta curry or pulses providing a similar amount of calories and protein;

** Vegetable rice or bread with salad providing a similar amount of calories and protein;

§ Mix vegetable curry or salad or any other vegetable providing a similar amount of calories and protein;

*** Peanuts or any other nuts providing a similar amount of calories and proteins

**** Carrot-halwa or pumpkin halwa or any other milk-based dessert providing a similar amount of calories and proteins



Note: 1 Cup = 250 ml; 1 plate pulao or dal or vegetables (100 g cooked food); 1 bowl fresh salad = (100 g);

1 serving peanuts = 1 ounce peanuts

Sample menu for lactating women

| Time | Food | Portion Size | Calories (Kcal) | Protein (g) |
|-----------|-------------------------------------|--------------|-----------------|-------------|
| Breakfast | Roti/toast/cereals* | 1 chapatti | 160 | 6 |
| | Egg fried‡ | 1 medium | 90 | 6 |
| | Milk† | 1 cup | 150 | 6 |
| Snack | Chicken sandwich‡ | 1 | 180 | 8 |
| | Apple or any other seasonal fruit ¶ | 1 medium | 80 | ----- |
| Lunch | Aloo qeema‡ | 1 plate | 222 | 11 |
| | Roti * | 2 chapatti | 320 | 12 |
| | Yogurt† | 1 cup | 150 | 6 |
| | Salad§ | 1 bowl§ | 25 | ----- |
| Snack | Shami kabab# | 1-2 | 90-180 | 6-12 |
| Dinner | Vegetable rice** | 1 plate | 187 | 4 |
| | Dal masoor | 1 plate | 140 | 8 |
| | Salad § | 1 bowl | 25 | ----- |
| | Raita | 1 cup | 150 | 6 |
| | Banana¶ | 1 medium | 80 | ----- |
| Snack | Peanuts §§ | 1 serving | 160 | 7 |
| | Carrot halwa*** | 1 cup | 250 | 6 |
| | Milk† | 1 cup | 150 | 6 |
| Total | | | 2609-2699 Kcal | 98-104 g |

*1 roti or 2 toast or ½ paratha or 1 cup fortified cereals providing a similar amount of calories and proteins; ‡ Egg fried or meat, chicken, fish or lentil or chickpea or red beans or any other pulses providing a similar amount of calories and proteins;

† 1 cup milk or 1 cup yogurt or 1 cup kheer or 1 ice cream or any other milk-based product providing a similar amount of calories and proteins;

¶ Apple or banana or any other seasonal fruit providing a similar amount of calories;

‡‡ Chicken sandwich or shami sandwich or egg sandwich or boiled chickpea or red beans or slice of pizza or peanut butter sandwich providing a similar amount of calories and proteins;

Shami kabab or red beans or boiled gram or meat based snack providing a similar amount of calories and proteins;

‡ Aloo qeema or mater qeema or beef curry or any other meat-based curry or beans or any other pulses providing a similar amount of calories and proteins;

**Vegetable rice or bread with salad providing a similar amount of calories and protein;

*** Carrot-halwa or pumpkin halwa or any other milk-based dessert providing a similar amount of calories and proteins;

§Salad or mix vegetable or any other vegetable providing a similar amount of calories;

§§ Peanuts or any other nuts providing a similar amount of calories and proteins; 8-10 glasses of water/liquid should be ensured for successful lactation

Note: 1 Cup = 250 ml; 1 plate pulao or dal or vegetables (100 g cooked food); 1 bowl fresh salad = (100 g);

1 serving peanuts = 1 ounce peanuts



Cost of healthy diets

The estimated cost based on the retail market price of food items of sample menus given above varies from Pak. Rupees (Rs.) 100-150 (\$1-1.50) for the whole menu per day. It is a fact that quality nutrient dense foods and diets will cost more than the monotonous unhealthy diets. However, consuming nutrient balanced diets including the recommended servings of milk and milk products, meat and pulses, cereals, vegetables and fruits, result in long-term benefits to consumers in terms of higher earnings and savings, improved health, improved physical productivity, reduced incidence of diseases and decreased medical costs.

Individuals with low income need to prioritize their household budgets to allocate a decent amount of money for food and re-appropriate expenditure towards purchase of healthy nutrient rich diets. The government may strengthen its social safety net programmes by providing food support in the form of food vouchers/coupons through an appropriate food distribution system, agriculture inputs support to the small-holders men and women farmers and link the farmers with markets and food distribution system.

Food may also be provided at subsidized rates to individuals living below the poverty line. Dietary guidelines can be used as an effective tool to promote nutrition literacy among individuals. Nutrition education and awareness plays a significant role in enabling individuals in general, and women in particular to make healthy, wise and appropriate food choices for themselves and their families.

DIETARY MESSAGES

Dietary messages have been developed in the light of scientific knowledge and epidemiological evidence to address nutritional and public health problems and to reduce and control the incidence of diseases and premature deaths. Efforts are made to make the dietary messages simple, easy, specific, comprehensible, meaningful, sensitive and culturally appropriate.

Maintain normal body weight by consuming all food groups and performing regular physical activity

Maintaining a normal (healthy) bodyweight through healthy dietary practices and regular exercise is vital for longevity and wellbeing, as both underweight and overweight are associated with increased prevalence of morbidity and mortality. Dietary diversification with regular exercise can minimize the adverse effects of obesity and overweight on health. Individuals who maintain normal body weight and are physically active have the lowest incidence of chronic disease and mortality.

Half of your daily cereals intake should include wholegrains

Whole grain cereals and pulses are good sources of providing energy, dietary fibre, proteins, carbohydrates, vitamins, minerals and antioxidants to the human body. They have been associated with reducing the risk of developing hypertension, metabolic syndrome, coronary heart disease, diabetes, cancers and other diseases.

Consumption of refined wheat and wheat products (pasta, noodles, white bread and other bakery products), rice and other cereals with low or negligible levels of dietary fibre have been linked to increased risk of digestive problems and colorectal cancers. About 45-65% of total calories shall come from carbohydrate containing foods⁹².

Eat five servings of fresh vegetables and fruits a day

Leafy green, deep yellow and red vegetables and fruits are rich sources of vitamin A, C, folic acid, iron, dietary fibre, potassium and antioxidants. They are free of fat and cholesterol, low in calories and contain many other beneficial compounds that prevent from overweight, obesity and incidence of chronic diseases and cancers. Five servings of vegetables and fruits per day are recommended for optimum health.

Take two to three servings of milk and milk products in a day

Milk and milk products are rich sources of calcium, phosphorous, magnesium, vitamin D, lactose, protein and fat, which are essential for skeletal growth and development. Adequate consumption of dairy products has been associated with strong bones and increased cognitive and physical outputs. For individuals who are overweight and obese or having cardiovascular diseases, skim milk is a preferred choice. Inadequate consumption of dairy products and lack of exposure to sunlight leads to rickets, osteomalacia and osteoporosis later in life. Two to three servings of milk and milk products per day are recommended to meet calcium and other essential nutrient requirements of the body.

Consume meat and meat products, fish and eggs in moderation

Meat and meat products, fish and eggs are rich in proteins, iron, zinc, vitamin A, vitamin B12 and integral component of hormones, enzymes and body immune system. Meat without fat (lean meat) is recommended over fatty meat (non-lean meat). Fish is recommended at least once in a week to supply protein and essential fatty acids to the body for growth and brain development. Excess amount of beef, sheep and goat meats shall be avoided to prevent high intake of saturated fat and cholesterol. Consumption of non-lean meat has been associated with increased cardiovascular diseases, breast and prostate cancers. Two to three servings of meat and meat products, chicken, fish and eggs per day are recommended to fulfill protein and other essential nutrient requirements of the body.

Encourage consumption of pulses to attain healthy growth

Pulses are rich in protein, dietary fibre, vitamins, minerals and essential amino acids particularly lysine, which complement the nutrients profile of wheat, rice and other cereals when consumed together. The protein content of pulses are almost equivalent to meat protein and therefore, could be used as a substitute where there is a shortage of meat or expensive to afford. They could be used as a preventive and therapeutic diets, as they tend to lower the risk of chronic non-communicable diseases i.e., hyperlipidemia, heart diseases, hypertension, diabetes and cancers.

Consume fortified flour, grains and their products

Fortified wheat flour and cereals are good sources of energy, proteins, vitamins and minerals. Use of fortified foods should be preferred over non-fortified foods to complement nutrients intake and combat macro and micronutrients deficiencies, improve health status and reduce risk of diseases. Micronutrient fortification has led to significant improvements in micronutrient status of women and children and reduction in prevalence of goiter, neural tube defects and other associated defects.

Limit consumption of edible oil and fat in cooking

Non-communicable diseases including diabetes, hypertension, cardiovascular diseases and cancers are reaching an epidemic proportion in Pakistan. These are responsible for the increasing burden on the health care system, human sufferings and premature deaths adversely affecting productivity and socio-economic development of the country. Consumption of saturated fat and trans-fatty acids increases low density lipoproteins (LDL) and cholesterol (bad fat) that increase the risk of hypertension and cardiovascular diseases. Evidence suggests that replacement of saturated fat and trans-fatty acids, with unsaturated cooking oil reduces bad fat (LDL & cholesterol) and increases good fat (high density lipoprotein) that decreases the risk of CVDs.

Reduce sugar intake, and limit intake of soft drinks, confectionaries, bakery products and commercial fruit drinks

Sugar sweetened drinks and foods with high sugar should be limited. Soft drinks, confectionaries, bakery products and commercial fruit drinks are energy dense foods and mostly provide empty calories increasing

the risk of overweight and obesity. These foods also contain large amount of sugar. Consumption of soft drinks and commercial sweet carbonated beverages has been strongly associated with increased weight gain, diabetes and pancreatic cancer. Consumption of complex carbohydrates is preferred over simple sugars. Reduction in sugar intake, soft drinks, confectionaries, bakery products, sweets and commercial sweetened fruit drinks is recommended.

Limit salt in cooking and always use iodized salt

In Pakistan, salt is more frequently used in diets, pickles and processed foods, and a potential risk factor for hypertension and other cardiovascular diseases. A positive association has been found between salt intake, hypertension, strokes and premature deaths. Salty foods are high in sodium. Sodium is one of the main dietary contributors to hypertension (high blood pressure). Even a small (1 g per person per day) reduction in salt intake will reduce deaths from strokes and heart attacks by more than 7% in each country. Avoid sprinkling of salt on fruits (oranges, lemons, grape fruits, guava, peaches, plums etc.) and beverages (soft drinks and orange, grape, peach, plum juices). Reduction in salt intake (< 5 g/day) is recommended for optimum health.

Limit consumption of fatty foods and highly processed foods

To improve health and nutritional status, substitution of junk (empty calories) and deep fried foods with healthy foods is essential. It is well established that consumption of junk and deep fried foods promote overweight and obesity that predispose children and adults to increased occurrence of diseases, resulting in impaired productivity and economic outputs. Highly processed foods, especially packaged, ready-made foods, are often high in fat, sugar and or salt and should be avoided. Instead eating raw vegetables and fruits and choosing fresh foods instead of packaged foods will contribute to maintain healthy diet.

Change sedentary lifestyle to physically active lifestyle

Physical activity is essential for skeletal growth and development, weight management, prevention of non-communicable diseases, increased productivity and better quality of life. At least half hour or more moderate exercise (e.g. walking) per day is recommended for physical fitness/health.

Exclusively breastfeed the baby in the first six months and continue breastfeeding along with complementary feeding at least for two years

Breast milk is a complete food for the first six months of life. It provides all the essential nutrients required by the baby for growth and development and therefore, no other liquid or food is required till the age of six months. After the age of six months, breastfeeding shall be complemented by nutrient dense and better varieties of safely prepared foods according to the infant and child's age. In order to allow child skin to form vitamin D, children over the age of six months should be exposed to sunlight in the early morning or late afternoon for few minutes a day or twice a week.

Women should increase intake of all the food groups daily, especially foods that are rich in iron and take extra care during pregnancy and lactation

Pregnancy and lactation impose extra demands of macro and micronutrients on women to support foetal growth and development, breastfeed the baby after birth and meet the nutritional requirements by eating varieties of safe and nutritious foods, especially foods that are rich in iron and other micronutrients.



Drink plenty of water each day

Drinking at least 8 glasses of clean and safe water per day is recommended for proper body functions. For a reference person of 65 kg, minimum required amount of drinking water is 2 liters (8 glasses) including all fluids. The requirement may vary according to the conditions (age, weather, work/activities, physiological status etc.).

Read nutrition labeling on packaged food products

Reading nutrition labels on packaged foods provide important information on the nutrients and food additives present in foods. Nutrition labeling facilitates the consumer to select the right and healthy food and enables to compare the nutrients composition of different brands of food products for the better selection.

ANNEXURE-I: Glycemic index and glycemic load of foods*

| Food | Glycemic Index | Carbohydrates | Glycemic Load |
|-----------------------------|----------------|---------------|---------------|
| Cereals | | | |
| White bread | 71 | 30 g | 10 |
| White rice | 89 | 150 g | 43 |
| Brown rice | 50 | 150 g | 16 |
| Barley (2/3 cup cooked) | 22 | 19 g | 4 |
| Sweet corn (2/3 cup cooked) | 47 | 20 g | 9 |
| Pasta (2/3 cup cooked) | 41 | 17 g | 7 |
| Spaghetti white | 58 | 180 g | 26 |
| Pulses | | | |
| Chana Dal (1/2 cup cooked) | 22 | 19 g | 4 |
| Lentils (1/2 cup cooked) | 21 | 15 g | 3 |
| Beans (1/2 cup cooked) | 30 | 15 g | 5 |
| Split Peas (1/2 cup cooked) | 25 | 13 g | 3 |
| Dairy Products | | | |
| Milk (1 cup) | 32 | 12 g | 4 |
| Yogurt (1 cup) | 19 | 14 g | 3 |
| Ice cream (1 cup) | 38 | 18 g | 7 |
| Vegetables | | | |
| Green peas | 51 | 80 g | 4 |
| Potato Boiled | 82 | 150 g | 21 |
| Fresh Fruits | | | |
| Bananas (1 medium) | 42 | 24 g | 10 |
| Apples (1 medium) | 37 | 21 g | 8 |
| Plums (2 whole) | 24 | 14 g | 3 |
| Peaches (1 large) | 28 | 14 g | 4 |
| Apricots (5 whole) | 34 | 15 g | 5 |
| Oranges (1 medium) | 40 | 13 g | 5 |
| Dried Fruits | | | |
| Apples (5 slices) | 29 | 15 g | 4 |
| Plum (3 pieces) | 29 | 15 g | 4 |
| Apricot (7 pieces) | 31 | 16 g | 5 |
| Peach (2 pieces) | 35 | 14 g | 5 |
| Pears (2 halves) | 29 | 24 g | 7 |

*Low glycemic foods. <http://optimalfoods.org/>

ANNEXURE-II: Personal hygiene and food safety guidelines*

- Keep yourself and kitchen surfaces neat and clean.
- Wash your hands, knives, equipment and cooking utensils before handling and use.
- Protect raw and cooked foods from flies, mosquitoes, insects and pests.
- Keep vegetables and fruits separate from raw meats and meat products.
- Use separate knives and chopping boards for vegetables, fruits and meat.
- Wash fruits and vegetables well, especially if eating raw.
- Use safe and clean water for cooking and drinking.
- Use fresh vegetable oil for cooking and frying.
- Avoid use of leftover vegetable oil for cooking and frying.
- Always purchase good quality packed and sealed vegetable oil from the market and avoid purchasing of unpacked oil from vendors.
- Cook food thoroughly, especially meat, eggs and fish, and store leftover foods in a safe place and at optimum temperatures.
- Refrigerate and freeze the food according to the recommended food safety guidelines.
- Reheat leftover foods thoroughly, until it is hot throughout before serving.
- Don't overcook vegetables to retain essential nutrients in the food.
- Use fresh food, don't use food that is old and spoiled.
- Keep food wastes in a container with cover to avoid contact of flies, mosquitoes, insects and pests and dispose it off quickly.
- Ban on use of pollutants, toxic chemicals, contaminated water and other industrial wastes in food processing and packing.
- Enhance the capacity of food inspectors, monitors and public health analysts to collect and analyze foods and beverages for pesticides, lead, arsenic and other toxic substances injurious to human health.
- Avoid meat exposure to open air and dust and take appropriate measures for its protection by installing wire gauze and cloth around the meat.



*<http://www.who.int/mediacentre/factsheets/fs399/en/>

ANNEXURE-III: Foods rich in omega fatty acids*

| Foods rich in omega-3 fatty acids (Alpha-linolenic acids) | Foods rich in omega-6 fatty acids (Linoleic acid) | Foods rich in omega-9 fatty acids (Oleic, Stearic acids and Erucic acids) |
|--|--|--|
| Salmon | Grains | Olives |
| Sardine | Nuts | Hazelnuts |
| Trout | Olive oil | Pecans |
| Tuna | Sunflower oil | Cashews |
| Mackerel | Safflower oil | Pistachios |
| Grains, Walnuts | | |

Oil and fats with different fatty acids composition

| Fats high in unsaturated fatty acids | Fats high in saturated fatty acids | Fats high in trans fatty acids |
|---|---|---------------------------------------|
| Corn | Coconuts | Margarine and vegetable ghee |
| Cotton seed | Red palm oil | Lard/cooking fat |
| Peanut | Butter | Deep fried foods |
| Soybean | Hydrogenated oil | Bakery products |
| Sunflower | Whole milk | |
| Olive | Fat from beef, mutton, chicken | |
| Sesame seed | Ghee | |
| Linseed | Lard | |
| Rapeseed | | |

Dietary Recommendations

- Energy intake from fat should range between 20-40% of total caloric intake.
- Energy intake from saturated fatty acids should not exceed 10% of the total energy intake.
- The concentration of trans fatty acids in the daily diet should not exceed 1% of the total energy intake from oils.
- Avoid or reduce consumption of deep-fried foods and fast foods.
- Avoid or reduce consumption of bakery products.
- Avoid or reduce consumption of Kababs prepared in lard and ghee.
- Avoid or reduce consumption of beef, mutton and chicken prepared in lard and ghee.
- Avoid or reduce the consumption of lard and ghee in daily cooking.
- Remove extra fat from meat and remove the skin from chicken.

*<http://www.livestrong.com/article/26035-list-omega-foods/>

ANNEXURE-IV: Recommended daily allowances for Pakistani population for selected major nutrients*

| Age (years) | Weight (kg) | Energy (kcal) | Protein (g) | Vitamin A (RE) | Vitamin D (ug) | Iron (mg) | Iodine (ug) | Zinc (mg) |
|--|-------------|---------------|-------------|----------------|----------------|-----------|-------------|-----------|
| <u>Children (boys and girls)</u> | | | | | | | | |
| 0-1 | 7.3 | 820 | 12 | 350 | 10 | 15 | 50 | 15 |
| 1-3 | 11.9 | 1250 | 23 | 400 | 10 | 20 | 90 | 15 |
| 3-5 | 15.9 | 1510 | 26 | 400 | 8.3 | 20 | 90 | 15 |
| 5-7 | 19.6 | 1710 | 30 | 400 | 4.2 | 20 | 90 | 15 |
| 7-10 | 25.9 | 1880 | 38 | 400 | 2.5 | 25 | 120 | 15 |
| <u>Children (boys)</u> | | | | | | | | |
| 10-12 | 34.0 | 2170 | 50 | 500 | 2.5 | 35 | 120 | 15 |
| 12-14 | 43.2 | 2360 | 64 | 600 | 2.5 | 35 | 150 | 15 |
| 14-16 | 54.5 | 2620 | 75 | 600 | 2.5 | 30 | 150 | 15 |
| 16-19 | 63.6 | 2820 | 84 | 600 | 2.5 | 30 | 150 | 15 |
| <u>Children (girls)</u> | | | | | | | | |
| 10-12 | 35.4 | 1925 | 52 | 500 | 2.5 | 30 | 120 | 15 |
| 12-14 | 44.2 | 2040 | 62 | 600 | 2.5 | 30 | 150 | 15 |
| 14-16 | 51.5 | 2135 | 69 | 550 | 2.5 | 20 | 150 | 15 |
| 16-19 | 54.6 | 2150 | 66 | 500 | 2.5 | 20 | 150 | 15 |
| Men moderately active – average 25 years | 55 | 2550 | 0.57 g/kg | 750 | 2.5 | 20 | 150 | 15 |
| Women moderately active – childbearing age | 46 | 2160 | 0.52 g/kg | 750 | 2.5 | 30 | 150 | 20 |
| Pregnant | ----- | +350 | +10 | 750 | 10 | 40 | 200 | 20 |
| Lactating | ----- | +350 | +26 | 1200 | 10 | 30 | 200 | 25 |

*Food Composition Table for Pakistan, Department of Agricultural Chemistry, University of Agriculture, Peshawar⁹³

ANNEXURE-V: Recommendations for infant and young child feeding*

- Initiate breast feeding within 1 hour of birth.
- Exclusive breast feeding for the first six months.
- Continue breast feeding until 2 years of age and beyond.
- Introduce complementary feeding at six months with small amounts of food and increase gradually as the child gets older.
- Gradually increase food consistency and variety.
- Be responsive to infant and child needs i.e., feed infant directly and assist older children.
- Feed child slowly and patiently and encourage them to eat but don't force.
- Practice good hygiene by washing hands frequently with soap particularly before preparing and eating foods and after using the toilet, use clean utensils and ensure proper food handling.
- Prepare complementary foods from locally available foods.
- Feed infants (aged 6-8 months) 2-3 times daily with soft foods which are easily digested by infants (banana, mashed potato, thick porridge, kheer, vegetable soup, kitchri) in addition to breast milk.
- Feed infant (aged 9-11 months) 3-4 times daily with soft foods which are easily digested by infants (banana, mashed potato, bread, thick porridge, kheer, egg, vegetable soup) in addition to breast milk.
- Feed child (aged 12-23 months) 3-4 times daily with homemade foods which are nutritious and digestible plus 1-2 additional snacks (bread, vegetables, egg, sandwich, banana, kheer, apple or any other fruit, egg, fish, chicken etc.) in addition to breast milk.
- Avoid giving sweets, candies, bakery products and beverages to children, and replace them with healthy foods such as fruits and vegetables, beans and pulses, nuts and dairy products.
- Serve child with fortified** cereals or vitamin-mineral supplements as needed.
- Increase fluid intake including more breast feeding and offer soft food to infants and children during illness.

*<http://www.who.int/mediacentre/factsheets/fs342/en/>

**GoP/WFP/NFA. Pakistan National Food Fortification Strategy 2017¹⁰⁵

ANNEXURE-VI: Complementary foods for infants > 6 months**RICE KHEER**

| Ingredients | Amount(g) |
|--|-----------|
| Rice | 60 |
| Milk | 30 |
| Nutritional Information per 100 g of edible portion | |
| Energy (Kcal) | 200 |
| Protein (g) | 11.4 |
| Fat/Oil (g) | 3.0 |
| Calcium (mg) | 150 |
| Phosphorus (mg) | 180 |
| Iron (mg) | 3 |

Preparation

Clean and wash rice and soak for ½ hr. Add ½ cup of water and rice in a deep saucepan. Cook on medium flame for 20-25 minutes till rice becomes soft. Mash rice with spoon or electric blender. Mix milk to it and again cook on medium flame for 10 minutes till it becomes soft and creamy in consistency and serve warm.



© FAO/Parvez Iqbal Paracha

KITCHRI

| Ingredients | Amount(g) |
|--|-----------|
| Rice | 60 |
| Dal Mung | 30 |
| Fat/Oil (g) | 5 |
| Nutritional Information per 100 g of edible portion | |
| Energy (Kcal) | 300 |
| Protein (g) | 11.4 |
| Fat/Oil (g) | 5.2 |
| Calcium (mg) | 120 |
| Phosphorus (mg) | 500 |
| Iron (mg) | 6 |

Preparation

Clean and wash rice and dal mung. Soak for ½ hour. Take one cup of water, rice, dal mung and rice in a deep saucepan. Cook on medium flame for 20-25 minutes till rice and dal mung become soft. Mash these and add one teaspoon of hot oil/fat, remove from the flame and serve warm.

SUJI KHEER

| Ingredients | Amount(g) |
|--|-----------|
| Suji | 20 |
| Milk | 120 |
| Nutritional Information per 100 g of edible portion | |
| Energy (Kcal) | 180 |
| Protein (g) | 5.2 |
| Fat/Oil (g) | 3.0 |
| Calcium (mg) | 150 |
| Phosphorus (mg) | 250 |
| Iron (mg) | 3 |

Preparation

Fry suji without oil in a deep saucepan till it becomes brown. Add milk to it and cook on medium flame for 10-15 minutes till it becomes soft and creamy in consistency. Turn off the flame and serve warm.

WHEAT DALIA

| Ingredients | Amount (g) |
|--|-------------------|
| Dalia (shredded wheat) | 20 |
| Milk | 120 |
| Water | 200 |
| Nutritional Information per 100 g of edible portion | |
| Energy (Kcal) | 380 |
| Protein (g) | 8.4 |
| Fat/Oil (g) | 6.0 |
| Calcium (mg) | 300 |
| Phosphorus (mg) | 380 |
| Iron (mg) | 2 |

Preparation

Thoroughly clean dalia and soak for ½ hr. Add 1 cup of water and dalia in a deep saucepan. Cook on low flame for 15-20 minutes till it becomes tender. Turn off the flame. Add milk and serve warm.

BANANA KHEER

| Ingredients | Amount (g) |
|--|-------------------|
| Banana | 80 |
| Milk | 120 |
| Nutritional Information per 100 g of edible portion | |
| Energy (Kcal) | 180 |
| Protein (g) | 3.0 |
| Fat/Oil (g) | 2.8 |
| Calcium (mg) | 160 |
| Phosphorus (mg) | 140 |
| Iron (mg) | 1.5 |

Preparation

Peel and mash banana with spoon. Add boiled milk. Mix to creamy consistency.

*Food Composition Table for Pakistan, Department of Agricultural Chemistry, University of Agriculture, Peshawar⁹³. The amount of sugar and salt are removed from the recipes as per WHO recommendations¹⁰².

ANNEXURE-VII: Food and portion sizes for different age groups

| Food Group | No. of Servings/day | | | | | Portion size/Exchanges |
|--|---------------------|------------|-------------|-------------|-----------|--|
| | 24-36 months | 3-10 years | 10-19 years | 19-60 years | >60 years | |
| Milk and milk products (Each Serving = 15 g carbohydrate, 6 g protein, 8 g fat, 150 calories) | 2 | 2-3 | 3-4 | 2-3 | 2-3 | Whole Milk= 1 cup Yogurt = 1 cup Cheese= 1 slice Kheer or Feerni= 1 cup Other milk-based products equivalent to nutrients supplied by 1 cup of milk |
| Cereals Each Serving = 30g carbohydrate, 6 g protein, 0-2 g fat, 160 calories | 1-2 | 2-4 | 5-6 | 4-5 | 4-5 | Bread (toast)= 2 slices Chapatti = 1 Cooked rice = 1 cup Cereals equivalent to nutrients supplied by 2 slices of bread |
| Vegetables Each serving = 5 g carbohydrates, 2 g protein and 25 calories Starchy Vegetables: Each serving =15 g carbohydrates, 3 g protein, 0-1 g Fat and 80calories | 1-2 | 1-2 | 2-3 | 2-3 | 2-3 | Cooked non-starchy vegetables=½ cup; Vegetables juice/soup= ½ cup Fresh vegetables/salad= 1 cup Starchy Vegetables Potato= 1(100 g) Maize = ½ cup Peas green= ½ cup |
| Fruits Each serving=20 g carbohydrates and 80 calories | 1-2 | 1-2 | 2-3 | 2-3 | 2-3 | Banana=1(medium size) or Apple= 1 or Peach=1 or Plums=2-3 or Apricots=3-4 |
| Meat and pulses • One serving of lean meat = 7 g protein, 3 g fat and 55 calories • One serving of meat with medium fat = 7 g protein, 5 g fat and 75 calories • One serving of meat with high fat =7 g protein, 8 g fat and 100 calories | 1-2 | 1-2 | 2-3 | 2-3 | 2-3 | Lean meat=28 g(2-3 small pieces) or Fish= 1-2 pieces or Egg=1 or Cooked pulses=½ cup |

ANNEXURE-VIII: Daily recommended intakes for energy and nutrients*

| Sex/Age | Body weight | Energy | Protein | Iron | Zinc | Vit. A | Vit. C | Folate |
|-------------------|-------------|--------|---------|---------|------|------------|--------|--------|
| Years | Kg | kcal | g | mg | mg | RE (µg) mg | | µg |
| Both sexes | | | | | | | | |
| 0-6 months | 6.0 | 524 | 11.6 | 0** | 1.1 | 375 | 25 | 80 |
| 6-11 months | 8.9 | 708 | 14.1 | 9 | 0.8 | 400 | 30 | 80 |
| 1-3 | 12.1 | 1022 | 14.0 | 6 | 8.4 | 400 | 30 | 160 |
| 4-6 | 18.2 | 1352 | 22.2 | 6 | 10.3 | 450 | 30 | 200 |
| 7-9 | 25.2 | 1698 | 25.2 | 9 | 11.3 | 500 | 35 | 300 |
| GIRLS | | | | | | | | |
| 10-17 | 46.7 | 2326 | 42.6 | 14 | 15.5 | 600 | 40 | 400 |
| Boys | | | | | | | | |
| 10-17 | 49.7 | 2824 | 47.8 | 17 | 19.2 | 600 | 40 | 400 |
| Women | | | | | | | | |
| | 55 | | | | | | | |
| 18-59 | | 2408 | 41.0 | 29 | 9.8 | 500 | 45 | 400 |
| Pregnant | | +278 | +6.0 | High*** | 15 | 800 | 55 | 600 |
| Breastfeeding | | +450 | +17.5 | 15 | 16.3 | 850 | 70 | 500 |
| 60 and over | | 2142 | 41.0 | 11 | 9.8 | 600 | 45 | 400 |
| Men | | | | | | | | |
| | 65 | | | | | | | |
| 18-59 | | 3091 | 49.0 | 14 | 14 | 600 | 45 | 400 |
| 60 and over | | 2496 | 49.0 | 14 | 14 | 600 | 45 | 400 |

*FAO¹⁰⁶. Family Nutrition Guide, Food and Agriculture Organization of the United Nations, Rome, 2004

** Full term babies are born with sufficient iron stores for first six months

*** Needs are so high that iron supplements are usually recommended for pregnant women

ANNEXURE-IX: Food as a source of nutrients

| Food | Constituents | Nutrients |
|------------------------|---|---|
| Cereals | Carbohydrates (starch, dietary fibre) | Energy, protein, vitamins (thiamine, riboflavin, niacin, vitamin E, folate), minerals (iron, zinc, phosphorous, magnesium) and fibre to reduce cholesterol, prevent constipation and diseases |
| Milk and milk products | Carbohydrates (lactose), proteins, fat, vitamins and minerals | Energy, proteins, vitamins (vitamin A , vitamin B12) minerals (calcium, phosphorous, magnesium) essential for bones, teeth, muscles and nerves conduction |
| Meat and pulses | Proteins, fat, vitamins and minerals | Proteins (amino acids), vitamins (Vitamin B12) and minerals (iron, zinc) |
| Vegetables | Vitamins, minerals, antioxidants, dietary fibre | Vitamins (B-carotene, B6, vitamin C, folic acid) minerals (iron, zinc, potassium) prevent constipation and diseases |
| Fruits | Vitamins, minerals, antioxidants, dietary fibre | Vitamins (B-carotene, B6, vitamin C, folic acid) minerals (iron, zinc, potassium) prevent constipation and diseases |
| Oil and fat | Essential and non-essential fatty acids | Energy, cholesterol and fatty acids |



ANNEXURE-X: Nutrients rich foods

Energy rich foods

- Oil, fat and sugar based foods (all desserts and halwa, confectionaries, bakery products)
- Fried and deep fried foods (Pakora, samosa, papar, fish, kabab, karhitikka)
- Fruits (Banana, apple, strawberry, melons, mangoes, sugar cane, pears, peaches, oranges, guava, honey and sugar)
- Starchy vegetables (potatoes)
- Cereals (wheat, maize, rice)
- Pulses (Chick pea roasted, boiled and cooked)
- Dried fruits and nuts (Apricots, figs, peaches, peanuts, walnuts, almonds)

Protein rich foods

- Beef, mutton, chicken, fish, liver, brain, milk
- Pulses (chick pea, red beans, lentil, mash and moong beans)
- Nuts (peanuts, almond, walnuts)
- Eggs
- Wheat, rice, maize

Iron rich foods

- Liver, beef, mutton
- Pulses
- Cereals
- Spinach and other dark green leafy vegetables
- Apple
- Fortified cereals

Zinc rich foods

- Beef, mutton, fish
- Pulses
- Cereals
- Fortified cereals



Calcium rich foods

- Milk and milk products
- Cabbage, broccoli and other dark green leafy vegetables
- Fortified cereals

Phosphorus rich foods

- Milk, yogurt, cheese
- Beef, mutton, chicken, salmon
- Pulses
- Nuts



Vitamin A rich foods

- Liver
- Colostrum and breast milk
- Dark green, yellow and red vegetables (spinach, pumpkin, carrots)
- Dark yellow and red fruits (water melon, mangoes, strawberries)
- Milk, yogurt, cheese
- Egg yolk

Folate rich foods

- Liver and kidney
- Dark green leafy vegetables
- Fortified cereals
- Egg yolk

Vitamin C rich foods

- Orange, lime, guava
- Dark green leafy vegetables
- Fresh milk

Sodium rich foods

- Table salt, baking powder
- Pickles, canned vegetables
- Canned processed foods, fast foods
- French fries, snacks, roasted and salted popcorns
- Roasted and salted nuts
- Salted cheese, instant soups
- Sauces and salad dressings

Potassium rich foods

- Banana, avocados,
- Potatoes, beans, spinach, mushrooms
- Dried fruits (apricots, peaches, raisins)
- Fish and yogurt

Trans rich foods

- Fried foods
- Bakery products
- Frosting in bakery products
- Margarine & industrially processed fats
- Processed foods

Sugar rich foods

- Bakery products
- Soft drinks and sugar added beverages
- Sugar cane juice and refined starchy foods

Salt rich foods

- Pickles
- Processed canned foods
- French fries and packaged snacks
- Salted nuts

ANNEXURE-XI: Healthy snacks

- All seasonal fruits
- Milk and milk products
- Fresh salad and green, yellow and red vegetables
- Peas and red beans
- Potato cutlets
- Shami kabab
- Shami sandwich with fresh salad
- Chicken sandwich with fresh salad
- Potato cutlet sandwich with fresh salad
- Cheese sandwich with fresh salad
- Egg sandwich with fresh salad
- Fruit chat
- Chana chat
- Dahi baray



ANNEXURE-XII: Recommendations on physical activity for different age groups*

Children and youth aged 5–17 years: Moderate-to-vigorous-intensity physical activity at least 60 minutes daily. The daily physical activity should preferably be aerobic and vigorous-intensity activities including muscle and bone strengthening exercises should be undertaken at least 3 times per week.

Adults aged 18-64 years: Moderate intensity physical activity (brisk walking, cycling, gardening, playing games, washing and cleaning clothes, car washing etc.) of 30 minutes at least five days a week or 15 minutes of vigorous intensity physical activity (weight lifting, running, bicycling, playing games, jumping ropes, heavy gardening etc.) at least five days a week or a combination of both if possible. Aerobic activity should be performed in bouts of at least 10 minutes duration. For additional health benefits, adults should increase their moderate-intensity aerobic physical activity to 300 minutes per week, or engage in 150 minutes of vigorous-intensity aerobic physical activity per week, or an equivalent combination of moderate- and vigorous-intensity activity. Muscle-strengthening activities should be done involving major muscle groups on 2 or more days a week.

Older adults: At least 30 minutes of moderate intensity aerobic physical activity daily or 15 minutes vigorous intensity aerobic physical activity daily or an equivalent combination of moderate to vigorous intensity activity are recommended. Older adults with poor mobility should perform physical activity to enhance their balance and prevents falls on 3 or more days per week. Muscle strengthening activities involving major muscles should also be done on 2 or more days a week. Older adults unable to perform moderate or vigorous intensity physical activity due to poor health, they should be as physically active as their abilities and conditions permits

*<http://www.who.int/mediacentre/factsheets/fs385/en/>.



ANNEXURE-XIII. Committees for Pakistan Dietary Guidelines for Better Nutrition**1. Steering/Core Committee**

| | |
|---|----------|
| 1. Dr. Mubarak Ali Ex-Member Food Security & Climate Change, Planning Commission | Chair |
| 2. Mr. Patrick T Evans, Ex-FAO Representative, Pakistan | Co-Chair |
| 3. Mr. M. Aslam Shaheen, Chief Nutrition, MPDR | Member |
| 4. Dr. Baseer Achakzi, Director Nutrition, MNHSRC | Member |
| 5. Dr. Shakil Ahmed Khan, Food Security Commissioner-II MFSR | Member |
| 6. Dr. M. Azeem Khan, Director General, NARC, Islamabad | Member |
| 7. Dr. Nomeena Anis, Nutritionist and Gender Focal Person FAO | Member |
| 8. Prof. Jehangir, K. Khalil, Ex-Dean, Agri. University, Peshawar | Member |
| 9. Dr. M. Sadiq Butt Dean NIFST University of Agriculture Faisalabad | Member |
| 10. Dr. Rezzan Khan, Consultant Clinical Dietitian Shifa Int. Hospital | Member |
| 11. Representative from WFP | Member |
| 12. Representative from UNICEF | Member |
| 13. Representative from WHO | Member |

2. Technical Committee

| | |
|---|--------|
| 1. Mr. M. Aslam Shaheen, Chief Nutrition, MPDR | Member |
| 2. Dr. Baseer Achakzai, Director Nutrition, MNHSRC | Member |
| 3. Dr. Rezzan Khan, Consultant Clinical Dietitian Shifa Int. Hospital | Member |
| 4. Dr. Nomeena Anis, Nutritionist and Gender Focal Person FAO | Member |
| 5. Prof. Ghazala Pervez Zaman, College of Home Economics, Lahore | Member |
| 6. Prof. Dr. Tahir Zahoor, NIFS&T, University of Agri. Faisalabad | Member |
| 7. Dr. Hajra Ahmad, Dept. of Home and Health Sci. AIOU, Islamabad | Member |
| 8. Dr. Mohammad Nasir, UVAS, Lahore | Member |
| 9. Dr. Nilofer Fatmi, Dow University of Health Sciences, Karachi | Member |
| 10. Dr. Romaina Iqbal, Aga Khan University, Karachi | Member |
| 11. Mr. Syed Saeed Qadir, Nutrition Officer, UNICEF, Islamabad | Member |

3. Author

Dr. Parvez Iqbal Paracha, Consultant Pakistan Dietary Guidelines for Better Nutrition, FAO, Islamabad

Co-Authors

1. Mr. M Aslam Shaheen, Chief Nutrition, MPDR
2. Dr. Nomeena Anis, Nutritionist and Gender Focal Person, FAO
3. Dr. Nazeer Ahmed, Deputy Chief Nutrition, MPDR
4. Ms. Nargis Hayat, Assistant Chief Nutrition, MPDR

4. List of Contributors

1. Dr. Aurangzeb, Director Nuclear Institute of Food and Agriculture (NIFA), Peshawar
2. Dr. Faqir Muhammad Anjum, Director, Institute of Home and Food Sciences Government College University, Faisalabad
3. Dr. Ghulam Sarwer Gilani, Health and Nutrition Advisor, Ottawa, Canada
4. Dr. S.M. Abid, Army Medical Corps, Lahore
5. Dr. Saeeda Raza, Director, Food Science and Product Development Institute, NARC, Islamabad
6. Dr. Iftikhar Alam, Chairman, Dept. Human Nutrition & Dietetics, Bacha Khan University Charsadda
7. Dr. Zia ud Din, Asstt. Prof. Dept. Human Nutrition, University of Agriculture, Peshawar
8. Dr. Sadaf Sardar, NPO SUN Secretariat, Islamabad
9. Mr. Bedar Abbas Ex-Deputy Chief Nutrition, MPDR

GLOSSARY

| | |
|---|--|
| Adolescence | A period between 10-19 years of age associated with rapid body growth and development, time to replenish body nutrients stores for future increased responsibilities |
| Acceptable Macronutrient Distribution Range | It is the percentage of total caloric intake that should come from a particular macronutrient (carbohydrates, protein and fat) each day |
| Anaemia | A state in which blood hemoglobin concentration falls below normal adversely affecting physical performance and productivity |
| Bakery products/confectionaries | All foods which are baked, fried or prepared from cereals (mainly wheat flour) in combination with sugar, oil and salt are called bakery products. They may include breads, biscuits, cakes, confectionaries (sweets), samosa, pakora and many others. They are rich in sugar, oil and salt and therefore, considered unhealthy. |
| Balanced diet | A diet which is composed of all food groups (grains, vegetables, fruit, meat and pulses, dairy products) and provide all the essential nutrients to the body in adequate amount |
| Basal metabolic rate | Amount of energy required to maintain metabolic functions of the human body |
| Clean and safe drinking water | Freshwater which is free from physical, microbial and chemical contaminants is called clean and safe drinking water |
| Contaminant | Any chemical, biological or any other substance that may make the food unsuitable for human consumption and possess potential health risk |
| Complementary feeding | A mashed food or thick porridge or soft semi-solid food introduced to the baby at the age of six months and continued till the child is on adult type diet in addition to breast milk |
| Exclusive breast feeding | A period from birth to six months of age when only breast milk is given to the baby |
| Fortified foods | Any food to which vitamins and minerals are added to increase the availability of nutrients to combat micronutrient deficiencies in population |
| Junk foods | Any food that provides empty calories and no other nutrients in substantial amount to meet the nutrients demand of the body. Fat and oil fried food, sugar based bakery products and confectionaries fall under the classification of junk foods |

| | |
|--------------------------|--|
| Malnutrition | A condition resulting from decreased or increased intake of foods. It includes both under-nutrition (underweight, wasting, stunting) and over-nutrition (overweight and obesity) |
| Macronutrient | Nutrients such as carbohydrates, proteins and fat which are required by the body in large amounts |
| Micronutrient | Nutrients such as vitamins and minerals which are required in small amounts |
| Nutrient reference value | Numerical values that are based on scientific data for purposes of nutrition labeling. |
| Obesity | A condition characterized by increased fatness. Body mass index greater than 30 indicates obesity in adults |
| Overweight | A condition characterized by higher weight than normal or reference mean weight. Body mass index between 25-29.9 indicates adults with overweight |
| Processed Foods | Food processing refers to a variety of processes carried out in food industry or at home and restaurants to make raw food and food ingredients into an edible form to improve its appearance, taste, color, shelf-life, digestibility and suitability for human consumption. Industrially produced processed foods are most commonly marketed in cans, card-board packages and boxes, nutritionally labeled and are ready to eat. |
| Snacks | Foods and beverages consumed/eaten between two big meals are called snacks. Snacks may be classified into energy dense (soft drinks, candies, bakery products, French fries etc.) and nutrients dense (milk, fresh fruits, dry nuts and fruits, salads, meat etc.). Healthy snacks help to maintain constant supply of nutrients and prevent body from hunger and overeating to reduce the risk of overweight, obesity and chronic diseases. |
| Stunting | Low height for age due to chronic energy deficiency |
| Wasting | Low weight for height due to acute energy deficiency |

REFERENCES

1. GOP. National Nutrition Survey 2011. Ministry of Planning, Development and Reform/Ministry of National Health Services, Regulations & Coordination, Government of Pakistan.
2. Jesmin, A., Yamamoto, SS., Malik, AA., Haque, A. Prevalence and determinants of chronic malnutrition among preschool children: A cross sectional study in Dhaka City, Bangladesh. *J Health Popul Nutr* 2011; 29:494-499.
3. Jamison, DT., Feachem, RG., Makgoba, MW., Bos, ER., Baingana, FK., Hofman, KJ., Rogo, KO. Disease and mortality in Sub-Saharan Africa. World Bank, 2006.
4. Tharakan, CT., Suchindran, CM. Determinants of child malnutrition-An intervention model for Botswana. *Nutr Res* 1999; 19:843-860.
5. Khan, AA., Bano, N., Salma, A. Child Malnutrition in South Asia: A comparative perspective. *South Asian Survey* 2007; 4:129-145.
6. World Bank. Repositioning nutrition as central to development: a strategy for large scale action: World Bank, Washington DC, 2006.
7. AICR. Cancer preventability estimates, Available at: <http://preventcancer.aicr.org/site/DocServer/UICCprWCD2011.pdf?docID=4781> [Accessed 10 Nov, 2016].
8. WHO. Effect of trans-fatty acid intake on blood lipids and lipoproteins. Available at: http://www.who.int/nutrition/publications/nutrientrequirements/tfa_systematic_review/en/ [Accessed 15 Dec, 2016].
9. WHO/MoH/Heartfile. National action plan for prevention and control of non-communicable diseases and health promotion in Pakistan, Islamabad, World Health Organization, Ministry of Health, Govt. of Pakistan and Heartfile, 2004.
10. IFPRI. Global Nutrition Report from promise to impact ending malnutrition by 2030. International Food Policy Research Institute, Washington, D.C. 2016.
11. Rafique, G., Azam, SI., White, F. Diabetes knowledge, beliefs and practice among people with diabetes attending a university hospital in Karachi, Pakistan. *Eastern Mediterranean Health Journal* 2006; 12:590-598.
12. George, A., Mensah, H., Brown, DW. An overview of cardiovascular disease burden in the United States. *Health Affairs* 2007; 26:38-48.
13. Yusuf, S., Hawken, S., Ounpuu, S., Dans, T., Avezum, A., Lanas, F., McQueen, M., Budaj, A, Pais, P., Varigos, J., Lisheng, L. Effect of potentially modifiable risk factors associated with myocardial infarction in 52 countries (the INTERHEART study): case-control study. *Lancet* 2004; 364:937-52.
14. WHO. Non-communicable diseases country profiles. World Health Organization, 2014.
15. UNICEF. Programming guide: Infant and young child feeding. Nutrition Section, United Nations Children's Fund, New York, 2011.
16. UNICEF. Breastfeeding. Available at: https://www.unicef.org/nutrition/index_24824.html [Accessed 2 Feb, 2017].

17. WHO. Indicators for assessing infant and young child feeding practices: Part III country profile. World Health Organization, Geneva, 2010.
18. WHO. Infant and young child feeding. Available at: <http://www.who.int/mediacentre/factsheets/fs342/en/> [Accessed 15 Nov, 2016].
19. WHO. Exclusive breastfeeding. Available at: http://www.who.int/nutrition/topics/exclusive_breastfeeding/en [Accessed 15 Nov, 2016].
20. UNICEF/WHO. Indicators for assessing infant and young child feeding practices: Part III Country Profile. World Health Organization, Geneva, 2010.
21. WHO. Child growth standards. Available at: http://www.who.int/childgrowth/standards/Technical_report.pdf [Accessed 20 Nov, 2016].
22. Shrimpton, R., Victora, CG., de Onis, Lima, RC., Blossner, M., Clugston, G. Worldwide timing of growth faltering: Implications for nutritional interventions. Available at: <http://www.pediatrics.org/cgi/content/full/107/5/e75> [Accessed 15 May, 2017].
23. Khan, MA., Rehman, Gn., Qazi, SA. Ari: Control of acute respiratory infections in Pakistan: Present status and future development. Federal ARI Cell, Children Hospital, PIMS, Pakistan 1990.
24. Saleemi, MA., Zaman, S., Akhtar, HZ., Jalil, F., Ashraf, N., Hanson, LA., Mellander, L. Feeding patterns, diarrhoeal illnesses and linear growth in 0-24 month-old children. *J Trop Paediatr* 2004; 50:164-169.
25. Musaiger, A.O., D'Souza, R. Anthropometric characteristics of Pakistani school children living in Bahrain. *J Immigrant Minority Health* 2009; 11-205.
26. Mushtaq, M.U., Gull, S., Khurshid, U., Shahid, U., Shad, M.A., Siddiqui, A.M. Prevalence and socio-demographic correlates of stunting and thinness among Pakistani primary school children. *BMC Public Health* 2011; 790-800.
27. Paracha, PI., Bakht, S., Paracha, SI., Vriesekoop, F., Khattak, IA., Zia ud Din, NiamatUllah. Nutritional status, dietary practices and physical activities of adolescents in public and private schools of Karachi, Pakistan. *Obes Res Open J* 2016; 3:30-39.
28. Aziz, S., Umme-e-Rubab, Noorulain, W., Majid, R., Hosain, K., Siddiqui, IA., Manzoor, S. Dietary patterns, height, weight centile and BMI of affluent school children and adolescents from three major cities of Pakistan. *J Coll Phys Surg Pak* 2010; 20:10-16.
29. Safdar, NF., Bertone-Johnson, E. Cordeiro, L., Jafar, TH., Cohen, NL. Dietary patterns of Pakistani adults and their association with sociodemographic, anthropometric and lifestyle factors. *J. Nutr Sci* 2013; 2:1-10.
30. WHO. Nutrition in adolescence: Issues in adolescent health and development, World Health Organization, Geneva, 2005.
31. Spear, B.A. Adolescent growth and development. *J Am Diet Assoc* 2002; 102(3):23-29.
32. Aziz, S., Noor-ul-Ain, W., Majeed, R., Khan, MA., Qayum, I., Ahmed, I., Hosain, K. Growth centile charts (anthropometric measurement) of Pakistani pediatric population. *J Pak Med Assoc* 2012; 62:367-377.

33. Hakeem, R., Sheikh, A.H., Asar, F. Assessment of linear growth of affluent urban Pakistani adolescents according to CDC 2000 references. Available at: <http://www.informaworld.com/smpp/title~content=t713723502~db=all~tab=issueslist~branchs=31-v31> [Accessed 10 Jan, 2017].
34. Paracha, PI., Waheed, H., Paracha, SI., Shahidullah, Bano, SS. Cardiovascular diseases in relation to anthropometric, biochemical and dietary intake in women: A case control study. *Obes Res Open J* 2015; 2:32-38.
35. Jafar, TH., Jafary, FH., Jessani, S., Chatury, N. Heart disease epidemic in Pakistan: Women and men at equal risk. *Am Heart J* 2005; 150:221-226.
36. GOP. National Health Survey (1990-994). Pakistan Medical Research Council, Government of Pakistan, Islamabad.
37. World Bank. Pakistan GDP per capita. Available at: <http://www.tradingeconomics.com/pakistan/gdp-per-capita> [Accessed 18 Nov, 2016].
38. Qidwai, W. Ageing Population: Status, challenges and opportunities for health care providers in Pakistan. *J Coll Phys Surg* 2009; 19:399-400.
39. Alam, I., Larbi, A., Pawelec, G., Paracha, PI. Relationship between anthropometric variables and nutrient intake in apparently healthy male elderly individuals: A study from Pakistan. *Nutr J* 2011; 10:111-119.
40. Lowe, NM., Ellahi, B., Bano, Q., Bangash, SA., Mitra, SR., Zaman, M. Dietary calcium intake, vitamin D status, and bone health in postmenopausal women in rural Pakistan. *J Health Popul Nutr* 2011; 29:465-470.
41. Qureshi, Z., Khan R. Diet intake trends among pregnant women in rural area of Rawalpindi, Pakistan. *J Ayub Med Coll Abbottabad* 2015; 27:684-88.
42. Henriksen, C., Brunvand, L., Stoltenberg, C., Trygg, K., Haug, E., Pedersen, JI. Diet and vitamin D status among pregnant Pakistani women in Oslo. *Eur J Clin Nutr* 1995; 49: 211-218.
43. Shaikh, F., Zeeshan, F., Hakeem, R., Basit, A., Fawwad, A., Hussain, A. Maternal dietary intake and anthropometric measurements of newborn at birth. *Open Diab J* 2014; 7:14-19.
44. Ijlal, F. Iron deficiency anaemia in the females and children of the Northern areas of Pakistan. Ph.D. thesis submitted to Quaid-I-Azam University, Islamabad, 2005.
45. Zia ud Din., Ambreen, S., Iqbal, Z., Iqbal, M., Ahmad, S., Hussain, M. Emmett, P. Inadequate dietary intake of women with antenatal psychological distress: A population based study in Pakistan. *J Food Nutr Res* 2014; 2:1021-1028.
46. Lassi, ZS., Mansoor, T., Salam, RA., Das, JK., Bhutta, ZA. Essential pre-pregnancy and pregnancy interventions for improved maternal, newborn and child health. *Reproductive Health* 2014; 11: (Suppl I)1-19.
47. Yakoob, MY., Menezes, EV., Soomro, T., Haws, RA., Darmstadt, GL., Bhutta, ZA. Reducing stillbirths behavioural and nutritional interventions before and during pregnancy. *BMC Pregnancy and Childbirth*. 2009; 9 (Suppl. 1):1-31.

48. IOM/NAP. Milk volume. Available at https://www.nap.edu/read/1577/chapter/6_ [Accessed 15 Nov, 2016].
49. Heymann, J., Raub, A., Earle, A. Breast feeding policy: a globally comparative analysis. Bull WHO 2013;91:398-406.
50. Dewey, KG., Finley, DA., Lonnerdal, B. Breast milk volume and composition during late lactation (7-20 months). J. Pediatr Gastroenterol Nutr 1984; 3:713-720.
51. Allen, LH. B vitamins in breast milk: Relative importance of maternal status and intake, and effects on infant status and function. Adv Nutr 2012; 3:362-369.
52. MPDR. Annual plan 2016, Ministry of Planning Development and Reform, Government of Pakistan.
53. FAO. Food balance sheets. Available at <http://www.fao.org/faostat/en/#data/FBS> [Accessed 16 Nov, 2016].
54. GOP. Chapter 11: Health and Nutrition, Pakistan Economic Survey, 2015-16. Ministry of Finance, Government of Pakistan, Islamabad. Available at: http://www.finance.gov.pk/survey/chapters_16/11_Health.pdf [Accessed 5 Nov, 2016].
55. GOP. Chapter 16:0 Poverty, Pakistan Economic Survey, 2015-16. Ministry of Finance, Government of Pakistan, Islamabad. Available at: http://www.finance.gov.pk/survey/chapters_16/Annexure_III_Poverty.pdf [Accessed 5 Nov, 2016].
56. Nazli, H., Shahzad, A., Mehmood, A. Food consumption patterns and nutritional status in Pakistan. Available at: <http://www.slideshare.net/m-hameed/food-consumption-patterns-and-nutritional-status-in-pakistan,2014> [Accessed 30 Oct, 2016].
57. FAO. Human energy requirements: Report of a joint FAO/WHO/UNU Expert Consultation. Food and Agriculture Organization, Rome, 2001.
58. FAO methodology for the measurement of food deprivation available at: http://www.fao.org/fileadmin/templates/ess/documents/food_security_statistics/metadata/Undernourishment_methodology.pdf [accessed 8 Feb, 2017].
59. FAO/IFAD/WFP. The state of food insecurity in the world 2015. Available at: <http://www.fao.org/3/a-i4646e.pdf> [Accessed 8 Feb, 2017].
60. WHO. Diet, nutrition and prevention of chronic diseases. Technical Report Series No. 916, World Health Organization, Geneva, 2003.
61. WHO. Policy statement and recommended actions to lower national salt intakes and lower death rates from high blood pressure and strokes in East Mediterranean Regions. World Health Organization, 2014.
62. WHO. Cancer prevention. Available at: <http://www.who.int/cancer/prevention/en/> [Accessed 4 Nov, 2016].
63. Malik, SJ. Food consumption patterns and nutrition. Available at: <http://www.slideshare.net/IFPRIDSG/s5-dr-sohail-j-malik-food-consumption-and-nutrition,2015> [Accessed 15 Sep, 2016].

64. WHO. Nutrient requirements and dietary guidelines. Available at <http://www.who.int/nutrition/publications/nutrient/en/> [Accessed 16 Nov, 2016].
65. Fast food industry of Pakistan growth and history. Available at: <http://ilm.com.pk/pakistan/pakistan-information/fast-food-industry-of-pakistan-growth-and-history/> [Accessed 2 Nov, 2016].
66. Glycemic index and diabetes. Available at: <http://www.diabetes.org/food-and-fitness/food/what-can-i-eat/understanding-carbohydrates/glycemic-index-and-diabetes.html> [Accessed 15 Dec, 2016].
67. Oh, K, Hu, FB., Cho, E., Rexrode, KM., Stampfer, MJ., Manson, J E., Liu, S., Willet, WC. Carbohydrate intake, glycemic index, glycemic load and dietary fiber in relation to risk of stroke in women. *Am J Epidemiol* 2005;16:161-169.
68. Greenwood, DC.,Threapleton, DE., Evans., CE. Glycemic index, glycemic load, carbohydrates, and type 2 diabetes: systematic review and dose-response meta-analysis of prospective studies. *Diabetes Care*. 2013; 36:4166-4171.63.
69. Food allergy. Available at <http://acaai.org/allergies/types/food-allergy> [Accessed 6 Jan 2017].
70. What is food allergy? Available at: <http://www.kidswithfoodallergies.org/page/what-is-a-food-allergy.aspx> [Accessed 15 Dec, 2016].
71. Processed food: Good or bad?<http://tribune.com.pk/story/671287/processed-food-good-or-bad/> [Accessed 15 Dec, 2016].
72. BMI classification. Available at: http://apps.who.int/bmi/index.jsp?introPage=intro_3.html [Accessed 13 Dec, 2016].
73. WHO. Waist circumference and waist-hip ratio. Report of a WHO expert consultation. Available at: http://apps.who.int/iris/bitstream/10665/44583/1/9789241501491_eng.pdf. [Accessed 12 Dec, 2016].
74. Imayama, I., Ulrich, CM., Alfano, CM., Wang, C., Xiao, L., Wener, MH. Effects of caloric restriction weight loss diet and exercise on inflammatory biomarkers in overweight/obese postmenopausal women: a randomized controlled trial. *Cancer Res* 2012; 79:2314-2326.
75. WHO. Global recommendations on physical activity for health. Available at: <http://www.who.int/dietphysicalactivity/publications/9789241599979/en/> [Accessed 15 Nov, 2016].
76. Donnelly, JE., Blair, SN., Jakicic, JM., Manore, MM., Rankin, JW., Smith, B. K. Appropriate physical activity intervention strategies for weight loss and prevention of weight regain for adults. *Med and Science in Sports and Exercise*, 2009; 41:459-471.
77. Dart, H., Nguyen, N., Colditz, G. A. Physical activity and chronic disease prevention: The young female athlete (pp. 163-179), Springer, 2016.
78. WHO. Physical inactivity a leading cause of disease and disability, warns WHO. Available at: <http://www.who.int/mediacentre/news/releases/release23/en/> [Accessed 15May, 2017].


79. Gilani Foundation. A 30 years of polling on eating habits of Pakistanis (1980-2010). Available at: <http://www.gilanifoundation.com/homepage/30years/30YearsOfPollingOnEatingHabitsOfPakistanis.pdf> [Accessed 15 May, 2017].
80. Stang, J., Story, M., Kossover, R. Promoting healthy eating and physical activity. Available at: http://www.epi.umn.edu/let/pubs/img/adol_ch6.pdf. [Accessed 20 Sep, 2016].
81. WHO. Tobacco. Available at: <http://www.who.int/mediacentre/factsheets/fs339/en/> [Accessed 5 Nov, 2016].
82. GOP. Pakistan Demographic and Health Survey (2012-013). National Institute of Population Studies, Government of Pakistan, Islamabad, Pakistan, 2013. Available at http://www.nips.org.pk/abstract_files/PDHS%20Final%20Report%20as%20of%20Jan%2022-2014.pdf [Accessed 10 Nov, 2016].
83. WHF. Totally avoidable risk factor for CVD. Available at: http://www.world-heart-federation.org/fileadmin/user_upload/documents/Fact_sheets/2012/Tobacco_avoidable_risk_factor_of_CVD.pdf [Accessed 20 Nov, 2016].
84. WHO. Food safety. Available at: <http://www.who.int/mediacentre/factsheets/fs399/en/>, 2015 [Accessed 5 Nov, 2016].
85. WHO. Water for health. Available at: http://www.who.int/water_sanitation_health/takingcharge.html [Accessed 5 Nov, 2016].
86. WHO. International standards for drinking water. Available at: <http://apps.who.int/iris/bitstream/10665/43845/1/a91160.pdf> [Accessed 4 Jan, 2017].
87. UNICEF. Available at: <http://nectar.com.pk/2016/09/10/unicef/> [Accessed 5 Nov, 2016].
88. Clean Drinking water for all (CDWA) project. Available at: http://www.environment.gov.pk/PRO_PDF/CDWA-Briefs.pdf [Accessed 5 Nov, 2016].
89. Memon, AN. Nutritional problems in Pakistan. Available at: <https://www.nuas.ac.jp/IHN/report/pdf/01/07.pdf> [Accessed 5 Nov, 2016].
90. WHO. Guidelines for drinking water quality. Available at: http://www.who.int/water_sanitation_health/dwq/fulltext.pdf [Accessed 13 Jan, 2017].
91. WHO/FAO. Preparation and use of food based dietary guidelines. Technical Report Series No. 880, World Health Organization, Geneva.
92. Eating well with Canada Food's Guide: A resource for educators and communicators. Available at: http://www.hc-sc.gc.ca/fn-an/alt_formats/hpfb-dgpsa/pdf/pubs/res-educat-eng.pdf [Accessed 15 Dec, 2016].
93. Food Composition Table for Pakistan (Revised 2001). Department of Agricultural Chemistry, University of Agriculture, Peshawar.
94. NIN/ICMR. Nutrient requirements and recommended dietary allowances for Indians. National Institute of Nutrition, Indian Council of Medical Research, Hyderabad, 2009.
95. WHO. Complementary feeding. Available at: http://www.who.int/nutrition/topics/complementary_feeding/en/ [Accessed 15 May, 2017].

96. WHO. Complementary feeding: Report of the global consultation. Available at http://www.who.int/nutrition/publications/Complementary_Feeding.pdf [Accessed 15 May, 2017].
97. GoP/UNICEF. Pakistan Infant and Young Child Feeding Strategy, 2016-20. Nutrition Wing, Ministry of National Health Services Regulations & Coordination, Government of Pakistan, 2015.
98. MeadJohnson. Toddler Nutrition. Available at: <https://www.meadjohnson.com/pediatrics/us-en/sites/hcp-usa/files/LB2264.pdf> [Accessed 15 May, 2017].
99. Kids Health. Nutrition guide for health. Available at: <http://Kidshealth.org/en/parents/toddler-food.html> [Accessed 15 May, 2017].
100. Growth reference data for 5-19 years. Available at: <http://www.who.int/growthref/en/> [Accessed 5 Nov, 2016]
101. IOM/NAP. Nutrition during pregnancy: part I Weight gain: Part II Nutrient supplements. Available at: <http://www.ncbi.nlm.nih.gov/books/NBK235235/> [Accessed 22 Oct, 2016].
102. WHO. Healthy diet fact sheet no. 394. Available at: http://www.who.int/elena/healthy_diet_fact_sheet_394.pdf [Accessed 5 Nov, 2016].
103. Venkaiah, K., Damayanti, K., Nayak, MU., Vijayaraghavan, K. Diet and nutritional status of rural adolescents in India. *Eur J Clin Nutr* 2002; 56:1119-1125.
104. Konje, JC., Ladipo, OA. Nutrition and obstructed labor. *Am J Clin Nutr* 2000; 72:291S-297S.
105. GoP/WFP/NFA. Pakistan National Food Fortification Strategy 2017. Ministry of National Health Services, Regulations and Coordination, Government of Pakistan.
106. FAO. Family nutrition guide. Food and Agriculture Organization of the United Nations, Rome, 2004.



Healthy People Strong Nation

Build healthy eating patterns



Drink plenty of water



Perform regular physical exercise



Adopt healthy life style and reduce stress



Limit Consumption of Unhealthy Foods



FAO Representation Office
NARC Premises, Park Road
Chak Shahzad, Islamabad
www.fao.org.pk

ISBN 978-92-5-131021-2



9 789251 310212

CA1868EN/1/10.18