

# Spectroscopic Methods in Organic Chemistry


## CHEM-6124, Organic Chemistry (Minor)

Online Lectures (MS)


Prof Dr Abdul Rauf Raza  
*Professor of Chemistry (Tenured)*  
Institute of Chemistry  
University of Sargodha, Sargodha

# Spectroscopy vs Spectrometry

1) Involves a physical change




2) a spectrum






1) Involves a chemical change



2) a spectrum



# Mass Spectrometry



# Energy of Accelerated Electrons


$$70 \text{ eV} = 70 \times (1.6 \times 10^{-19}) = 1.12 \times 10^{-17} \text{ J / partical} \times (6.02 \times 10^{23}) \\ = 6.75 \times 10^6 \text{ J . mol}^{-1} = 6.75 \times 10^3 \text{ kJ . mol}^{-1}$$

$$\text{since } E = h\nu \quad \text{so } \nu = E/h = \frac{1.12 \times 10^{-17} \text{ J}}{6.63 \times 10^{-34} \text{ J.s}} = 1.69 \times 10^{16} \text{ s}^{-1} (\text{Hz})$$

$$\text{since } c = \nu \lambda \quad \text{so } \lambda = c/\nu = \frac{3 \times 10^8 \text{ m.s}^{-1}}{1.69 \times 10^{16} \text{ s}^{-1}} = 1.78 \times 10^{-8} \text{ m}$$

$$\lambda = 17.8 \text{ nm} = 178 \text{ \AA}^\circ (\text{VUV: } 3 - 200 \text{ nm})$$


# A Mass Spectrometer




Prof Dr Abdul Rauf Raza, University of Sargodha

# Ionization Techniques







Techniques	MM (amu)
Electron Impact (EI)	1,000
Chemical Ionization (CI)	1,000
Atmospheric pressure	2,000
Chemical Ionization (APCI)	
Fast Atomic Bombardment (FAB)	6,000
Electro Spray Ionization (ESI)	10,000
Matrix Assisted Laser	
Desorption Ionization (MALDI)	500,000




# Electron Impact (EI)



# Chemical Ionization (CI)




# APCI / ESI



## ESI:



- 1) Ionization takes place first rather vaporization
- 2) Analyte may be an organic molecule or a polymer/biopolymer
- 3) Matrix is a buffer of  $\text{K}^+$ ,  $\text{Na}^+$ ,  $^+\text{NH}_4$  ions
- 4) Molecular ions  $[\text{M}+\text{H}]$ ,  $[\text{M}+\text{Na}]$ ,  $[\text{M}+\text{K}]$ ,  $[\text{M}+\text{NH}_4]$

# Fast Atomic Bombardment (FAB)




# MALDI

Matrix Assisted Laser Desorption Ionization




Matrix ( $\lambda_{max}$ )




# LASER Principle

Light Amplification by Stimulated Emission of Radiations




$$E_2 - E_1 = \Delta E = h\nu$$

# LASER Principle



# Fragmentation



$$\text{K.E.} = \text{Ve} \quad \Rightarrow \quad mv^2/2 = \text{Ve}$$

$$v^2 = 2 \text{ Ve} / m \dots\dots \text{eq 1}$$

since  $F_c \propto H \nu$

$$H^2 e^2 r^2 / m^2 = 2 \text{ Ve} / m$$



$$mv^2/r = He\nu \quad v^2/\nu = He r/m$$

$$2V/H^2 r^2 = e/m$$


$$v^2 = H^2 e^2 r^2 / m^2 \dots\dots \text{eq 2}$$

$$m/e = H^2 r^2 / 2V$$


# $\alpha$ -Cleavage




# $\alpha$ -Cleavage




# Fragmentation ( $\alpha$ -Cleavage)




# Stability of Carbocation




# $\beta$ -Ketoacid Rearrangement



# McLafferty ( $\gamma$ -H) Rearrangement



# *Retro* Diels-Alder



# Isotopic Abundances of Cl

Number of [M] in MS spectrum =  $n + 1$ ; where  $n$  = number of halogens

Ratio of lines in MS spectrum =  $(a + b)^n$ ; where  $n$  = number of halogens

$$\begin{aligned}(a + b)^1 &= 0.75a + 0.25b \\ &= 3a + 1b\end{aligned}$$

$$\begin{aligned}(a + b)^2 &= a^2 + 2ab + b^2 \\ &= (0.75)^2 a^2 + 2(0.75)a (0.25)b + (0.25)^2 b^2 \\ &= (0.56)a^2 + (0.38)ab + (0.06)b^2 \\ &= 9a^2 + 6ab + 1b^2\end{aligned}$$

$$\begin{aligned}(a + b)^3 &= a^3 + 3a^2b + 3ab^2 + b^3 \\ &= (0.75)^3 a^3 + 3(0.75)^2 a^2 (0.25)b + 3(0.75)a (0.25)^2 b^2 + (0.25)^3 b^3 \\ &= (0.42)a^3 + (0.42)a^2b + (0.14)ab^2 + (0.016)b^3 \\ &= 30a^3 + 30a^2b + 10ab^2 + 1b^3\end{aligned}$$


# Isotopic Abundances of Br

$$(a + b)^1 = 0.5a + 0.5b \\ = 1a + 1b$$

$$(a + b)^2 = a^2 + 2ab + b^2 \\ = (0.5)^2 a^2 + 2(0.5)a(0.5)b + (0.5)^2 b^2 \\ = (0.25)a^2 + (0.5)ab + (0.25)b^2 \\ = 1a^2 + 2ab + 1b^2$$



$$(a + b)^3 = a^3 + 3a^2b + 3ab^2 + b^3 \\ = (0.50)^3 a^3 + 3(0.50)^2 a^2 (0.50)b + 3(0.50)a (0.50)^2 b^2 + (0.50)^3 b^3 \\ = (0.125)a^3 + (0.375)a^2b + (0.375)ab^2 + (0.125)b^3 \\ = 1a^3 + 3a^2b + 3ab^2 + 1b^3$$

# GCMS




# GC EIMS

ANHINDI #511 RT: 12.04 AC: 1 NL: 7.04 EM  
T: [0,0] + c Full ms (50.00-600.00)




$C_{13}H_{14}O_2$   
 $m/z = 202.1$  amu

# Fragmentation




# EIMS





Prof Dr Abdul Rauf Raza, University of  
Sargodha

# EIMS





# EIMS

FA-19-26 #687 RT: 15.15 AV: 1 NL: 4.40E7  
T: [0,0] + c Full ms [50.00-900.00]



$m/z = 319, 321$  amu (3:1)


# EIMS




$m/z = 353, 355, 357$  amu (9:6:1)

# High Resolution ESIMS


SA-11-45\_Pos\_full #1-35 RT: 0.00-0.50 AV: 35 NL: 1.10E8  
T: FTMS + p NSI Full ms [100.00-2000.00]



# High Resolution ESIMS




# High Resolution ESIMS



Chemical Formula:  $C_{30}H_{25}ClN_2O_3^+$

Exact mass: 497.163 amu


# High Resolution ESIMS



Chemical Formula:  $C_{30}H_{25}BrN_2O_3^+$

Exact mass: 541.112 amu

# High Resolution ESIMS



Chemical Formula:  $C_{30}H_{26}Cl_2N_2O_3^+$

Exact mass: 532.13 amu