

FIGURE 2.21 Different right-hand and left-hand limits at the origin.

In this section we extend the limit concept to *one-sided limits*, which are limits as  $x$  approaches the number  $s_0$  from the left-hand side (where  $x < s_0$ ) or the right-hand side ( $x > s_0$ ) only. We also analyze the graphs of certain rational functions as well as other functions with limit behavior as  $x \rightarrow \pm\infty$ .

### One-Sided Limits

To have a limit  $L$  as  $x$  approaches  $c$ , a function  $f$  must be defined on *both sides* of  $c$  and its values  $f(x)$  must approach  $L$  as  $x$  approaches  $c$  from either side. Because of this, ordinary limits are called **two-sided**.

If  $f$  fails to have a two-sided limit at  $c$ , it may still have a one-sided limit, that is, a limit if the approach is only from one side. If the approach is from the right, the limit is a **right-hand limit**. From the left, it is a **left-hand limit**.

The function  $f(x) = x/|x|$  (Figure 2.21) has limit 1 as  $x$  approaches 0 from the right, and limit  $-1$  as  $x$  approaches 0 from the left. Since these one-sided limits are different, there is no single number that  $f(x)$  approaches as  $x$  approaches 0. We say that  $f$  has a **two-sided limit** at 0.

Intuitively, if  $f(x)$  is defined on an interval  $(c, b)$ , we say that  $f$  has a **right-hand limit** at  $c$  if  $f(x)$  approaches  $L$  as  $x$  approaches  $c$  from within that interval. We write

$$\lim_{x \rightarrow c^+} f(x) = L.$$

The symbol " $x \rightarrow c^+$ " means that we consider only values of  $x$  greater than  $c$ .

Similarly, if  $f(x)$  is defined on an interval  $(a, c)$ , where  $a < c$  and  $x$  approaches arbitrarily close to  $M$  as  $x$  approaches  $c$  from within that interval, then  $f$  has **left-hand limit**  $M$  at  $c$ . We write

$$\lim_{x \rightarrow c^-} f(x) = M.$$

The symbol " $x \rightarrow c^-$ " means that we consider only  $x$  values less than  $c$ .

These informal definitions are illustrated in Figure 2.22. For the function  $f(x) = x/|x|$  in Figure 2.21 we have

$$\lim_{x \rightarrow 0^+} f(x) = 1 \quad \text{and} \quad \lim_{x \rightarrow 0^-} f(x) = -1.$$




FIGURE 2.22 (a) Right-hand limit as  $x$  approaches  $c$ . (b) Left-hand limit as  $x$  approaches  $c$ .




FIGURE 2.23  $\lim_{x \rightarrow -2^+} \sqrt{4-x^2} = 0$  and  $\lim_{x \rightarrow 2^-} \sqrt{4-x^2} = 0$  (Example 1)

### EXAMPLE 1 One-Sided Limits for a Semicircle

The domain of  $f(x) = \sqrt{4-x^2}$  is  $[-2, 2]$ ; its graph is the semicircle in Figure 2.23. We have

$$\lim_{x \rightarrow -2^+} \sqrt{4-x^2} = 0 \quad \text{and} \quad \lim_{x \rightarrow 2^-} \sqrt{4-x^2} = 0.$$

The function does not have a left-hand limit at  $x = -2$  or a right-hand limit at  $x = 2$ . It does not have ordinary two-sided limits at either  $-2$  or  $2$ .

One-sided limits have all the properties listed in Theorem 1 in Section 2.2. The right-hand limit of the sum of two functions is the sum of their right-hand limits, and so on. The theorems for limits of polynomials and rational functions hold with one-sided limits, as does the Sandwich Theorem and Theorem 5. One-sided limits are related to limits in the following way.

### THEOREM 6

A function  $f(x)$  has a limit as  $x$  approaches  $c$  if and only if it has left-hand and right-hand limits there and these one-sided limits are equal.

$$\lim_{x \rightarrow c} f(x) = L \iff \lim_{x \rightarrow c^-} f(x) = L \quad \text{and} \quad \lim_{x \rightarrow c^+} f(x) = L.$$




FIGURE 2.24 Graph of the function in Example 2

### EXAMPLE 2 Limits of the Function Graphed in Figure 2.24

At  $x = 0$ :  $\lim_{x \rightarrow 0^+} f(x) = 1$ ,  
 $\lim_{x \rightarrow 0^-} f(x)$  and  $\lim_{x \rightarrow 0} f(x)$  do not exist. The function is not defined to the left of  $x = 0$ .

At  $x = 1$ :  $\lim_{x \rightarrow 1^-} f(x) = 0$  even though  $f(1) = 1$ ,  
 $\lim_{x \rightarrow 1^+} f(x) = 1$ ,  
 $\lim_{x \rightarrow 1} f(x)$  does not exist. The right- and left-hand limits are not equal.

At  $x = 2$ :  $\lim_{x \rightarrow 2^-} f(x) = 1$ ,  
 $\lim_{x \rightarrow 2^+} f(x) = 1$ ,  
 $\lim_{x \rightarrow 2} f(x) = 1$  even though  $f(2) = 2$ .

At  $x = 3$ :  $\lim_{x \rightarrow 3^-} f(x) = \lim_{x \rightarrow 3^+} f(x) = \lim_{x \rightarrow 3} f(x) = f(3) = 2$ .

At  $x = 4$ :  $\lim_{x \rightarrow 4^-} f(x) = 1$  even though  $f(4) \neq 1$ ,  
 $\lim_{x \rightarrow 4^+} f(x)$  and  $\lim_{x \rightarrow 4} f(x)$  do not exist. The function is not defined to the right of  $x = 4$ .

At every other point  $c$  in  $[0, 4]$ ,  $f(x)$  has limit  $f(c)$ .

### Precise Definitions of One-Sided Limits

The formal definition of the limit in Section 2.3 is readily modified for one-sided limits.

121 / 1564






FIGURE 2.25 Intervals associated with the definition of right-hand limit.




FIGURE 2.26 Intervals associated with the definition of left-hand limit.




FIGURE 2.27  $\lim_{x \rightarrow 0^+} \sqrt{x} = 0$  in Example 3

**DEFINITIONS** Right-Hand, Left-Hand Limits

We say that  $f(x)$  has **right-hand limit**  $L$  at  $x_0$ , and write

$$\lim_{x \rightarrow x_0^+} f(x) = L \quad (\text{See Figure 2.25})$$

if for every number  $\epsilon > 0$  there exists a corresponding number  $\delta > 0$  such that for all  $x$

$$x_0 < x < x_0 + \delta \implies |f(x) - L| < \epsilon.$$

We say that  $f$  has **left-hand limit**  $L$  at  $x_0$ , and write

$$\lim_{x \rightarrow x_0^-} f(x) = L \quad (\text{See Figure 2.26})$$

if for every number  $\epsilon > 0$  there exists a corresponding number  $\delta > 0$  such that for all  $x$

$$x_0 - \delta < x < x_0 \implies |f(x) - L| < \epsilon.$$

**EXAMPLE 3** Applying the Definition to Find Delta

Prove that

$$\lim_{x \rightarrow 0^+} \sqrt{x} = 0.$$

**Solution** Let  $\epsilon > 0$  be given. Here  $x_0 = 0$  and  $L = 0$ , so we want to find a  $\delta > 0$  such that for all  $x$

$$0 < x < \delta \implies |\sqrt{x} - 0| < \epsilon.$$

or

$$0 < x < \delta \implies \sqrt{x} < \epsilon.$$

Squaring both sides of this last inequality gives

$$x < \epsilon^2 \quad \text{if} \quad 0 < x < \delta.$$

If we choose  $\delta = \epsilon^2$  we have

$$0 < x < \delta = \epsilon^2 \implies \sqrt{x} < \epsilon.$$

or

$$0 < x < \epsilon^2 \implies |\sqrt{x} - 0| < \epsilon.$$

According to the definition, this shows that  $\lim_{x \rightarrow 0^+} \sqrt{x} = 0$  (Figure 2.27). ■

The functions examined so far have had some kind of limit at each point of interest. In general, that need not be the case.

**EXAMPLE 4** A Function Oscillating Too Much

Show that  $y = \sin(1/x)$  has no limit as  $x$  approaches zero from either side (Figure 2.28).

**Solution** As  $x$  approaches zero, its reciprocal,  $1/x$ , grows without bound and the values of  $\sin(1/x)$  cycle repeatedly from  $-1$  to  $1$ . There is no single number  $L$  that the function's




FIGURE 2.28 The function  $y = \sin(1/x)$  has neither a right-hand nor a left-hand limit as  $x$  approaches zero (Example 4).

values stay increasingly close to  $L$  as  $x$  approaches zero. This is true even if we restrict  $x$  to positive values or to negative values. The function has neither a right-hand limit nor a left-hand limit at  $x = 0$ . ■

**Limits Involving  $(\sin \theta)/\theta$**

A central fact about  $(\sin \theta)/\theta$  is that in radian measure its limit as  $\theta \rightarrow 0$  is 1. We can see this in Figure 2.29 and confirm it algebraically using the Sandwich Theorem.




FIGURE 2.29 The graph of  $f(\theta) = (\sin \theta)/\theta$ .




FIGURE 2.30 The figure for the proof of Theorem 7.  $PA/OA = \tan \theta$ , but  $OA = 1$ , so  $PA = \tan \theta$ .

**THEOREM 7**

$$\lim_{\theta \rightarrow 0} \frac{\sin \theta}{\theta} = 1 \quad (\theta \text{ in radians}) \quad (1)$$

**Proof** The plan is to show that the right-hand and left-hand limits are both 1. Then we will know that the two-sided limit is 1 as well.

To show that the right-hand limit is 1, we begin with positive values of  $\theta$  less than  $\pi/2$  (Figure 2.30). Notice that

$$\text{Area } \triangle OAP < \text{area sector } OAP < \text{area } \triangle OAT.$$

(b) Equation (1) does not apply to the original fraction. We need a  $2x$  in the denominator, not a  $5x$ . We produce it by multiplying numerator and denominator by  $2/5$ .

$$\begin{aligned} \lim_{x \rightarrow 0} \frac{\sin 2x}{5x} &= \lim_{x \rightarrow 0} \frac{(2/5) \cdot \sin 2x}{(2/5) \cdot 5x} \\ &= \frac{2}{5} \lim_{x \rightarrow 0} \frac{\sin 2x}{2x} \quad \text{Now Eq. (1) applies with } \theta = 2x. \\ &= \frac{2}{5} (1) = \frac{2}{5} \end{aligned}$$




FIGURE 2.31 The graph of  $y = 1/x$ .

**Finite Limits as  $x \rightarrow \pm\infty$**

The symbol for infinity ( $\infty$ ) does not represent a real number. We use  $\infty$  to describe the behavior of a function when the values in its domain or range outgrow all finite bounds. For example, the function  $f(x) = 1/x$  is defined for all  $x \neq 0$  (Figure 2.31). When  $x$  is positive and becomes increasingly large,  $1/x$  becomes increasingly small. When  $x$  is negative and its magnitude becomes increasingly large,  $1/x$  again becomes small. We summarize these observations by saying that  $f(x) = 1/x$  has limit 0 as  $x \rightarrow \pm\infty$ , or that 0 is a limit of  $f(x) = 1/x$  at infinity and negative infinity. Here is a precise definition.

**DEFINITIONS** Limit as  $x$  approaches  $\infty$  or  $-\infty$

1. We say that  $f(x)$  has the **limit  $L$  as  $x$  approaches infinity** and write

$$\lim_{x \rightarrow \infty} f(x) = L$$

if, for every number  $\epsilon > 0$ , there exists a corresponding number  $M$  such that for all  $x$

$$x > M \implies |f(x) - L| < \epsilon.$$

2. We say that  $f(x)$  has the **limit  $L$  as  $x$  approaches minus infinity** and write

$$\lim_{x \rightarrow -\infty} f(x) = L$$

if, for every number  $\epsilon > 0$ , there exists a corresponding number  $N$  such that for all  $x$

$$x < N \implies |f(x) - L| < \epsilon.$$

Intuitively,  $\lim_{x \rightarrow \infty} f(x) = L$  if, as  $x$  moves increasingly far from the origin in the positive direction,  $f(x)$  gets arbitrarily close to  $L$ . Similarly,  $\lim_{x \rightarrow -\infty} f(x) = L$  if, as  $x$  moves increasingly far from the origin in the negative direction,  $f(x)$  gets arbitrarily close to  $L$ .

The strategy for calculating limits of functions as  $x \rightarrow \pm\infty$  is similar to the one for finite limits in Section 2.2. There we first found the limits of the constant and identity functions  $y = k$  and  $y = x$ . We then extended these results to other functions by applying a theorem about limits of algebraic combinations. Here we do the same thing, except that the starting functions are  $y = k$  and  $y = 1/x$  instead of  $y = k$  and  $y = x$ .




FIGURE 2.32 The geometry behind the argument in Example 6.

The basic facts to be verified by applying the formal definition are

$$\lim_{x \rightarrow \pm\infty} k = k \quad \text{and} \quad \lim_{x \rightarrow \pm\infty} \frac{1}{x} = 0. \quad (3)$$

We prove the latter and leave the former to Exercises 71 and 72.

**EXAMPLE 6** Limits at Infinity for  $f(x) = \frac{1}{x}$

Show that

- (a)  $\lim_{x \rightarrow \infty} \frac{1}{x} = 0$                       (b)  $\lim_{x \rightarrow -\infty} \frac{1}{x} = 0$ .

**Solution**

(a) Let  $\epsilon > 0$  be given. We must find a number  $M$  such that for all  $x$

$$x > M \implies \left| \frac{1}{x} - 0 \right| = \left| \frac{1}{x} \right| < \epsilon.$$

The implication will hold if  $M = 1/\epsilon$  or any larger positive number (Figure 2.32). This proves  $\lim_{x \rightarrow \infty} (1/x) = 0$ .

(b) Let  $\epsilon > 0$  be given. We must find a number  $N$  such that for all  $x$

$$x < N \implies \left| \frac{1}{x} - 0 \right| = \left| \frac{1}{x} \right| < \epsilon.$$

The implication will hold if  $N = -1/\epsilon$  or any number less than  $-1/\epsilon$  (Figure 2.32). This proves  $\lim_{x \rightarrow -\infty} (1/x) = 0$ .

Limits at infinity have properties similar to those of finite limits.

**THEOREM 3** Limit Laws as  $x \rightarrow \pm\infty$

If  $L$ ,  $M$ , and  $k$  are real numbers and

$$\lim_{x \rightarrow \pm\infty} f(x) = L \quad \text{and} \quad \lim_{x \rightarrow \pm\infty} g(x) = M,$$

then


- Sum Rule:**  $\lim_{x \rightarrow \pm\infty} (f(x) + g(x)) = L + M$
- Difference Rule:**  $\lim_{x \rightarrow \pm\infty} (f(x) - g(x)) = L - M$
- Product Rule:**  $\lim_{x \rightarrow \pm\infty} (f(x) \cdot g(x)) = L \cdot M$
- Constant Multiple Rule:**  $\lim_{x \rightarrow \pm\infty} (k \cdot f(x)) = k \cdot L$
- Quotient Rule:**  $\lim_{x \rightarrow \pm\infty} \frac{f(x)}{g(x)} = \frac{L}{M}, \quad M \neq 0$
- Power Rule:** If  $r$  and  $s$  are integers with no common factors,  $s \neq 0$ , then  $\lim_{x \rightarrow \pm\infty} (f(x))^{r/s} = L^{r/s}$  provided that  $L^{r/s}$  is a real number. (If  $s$  is even, we assume that  $L > 0$ .)

These properties are just like the properties in Theorem 1, Section 2.2, and we use them the same way.




**EXAMPLE 7** Using Theorem 8

$$\begin{aligned} \text{(a)} \quad \lim_{x \rightarrow \infty} \left( 5 + \frac{1}{x} \right) &= \lim_{x \rightarrow \infty} 5 + \lim_{x \rightarrow \infty} \frac{1}{x} && \text{Sum Rule} \\ &= 5 + 0 = 5 && \text{Known limits} \\ \text{(b)} \quad \lim_{x \rightarrow \infty} \frac{\pi\sqrt{x}}{x^2} &= \lim_{x \rightarrow \infty} \pi\sqrt{x} \cdot \frac{1}{x^2} \\ &= \lim_{x \rightarrow \infty} \pi\sqrt{x} \cdot \lim_{x \rightarrow \infty} \frac{1}{x^2} && \text{Product rule} \\ &= \pi\sqrt{3} \cdot 0 = 0 && \text{Known limits} \end{aligned}$$



**FIGURE 2.33** The graph of the function in Example 8. The graph approaches the line  $y = 5/3$  as  $|x|$  increases.



**FIGURE 2.34** The graph of the function in Example 9. The graph approaches the  $x$ -axis as  $|x|$  increases.

**Limits at Infinity of Rational Functions**

To determine the limit of a rational function as  $x \rightarrow \pm\infty$ , we can divide the numerator and denominator by the highest power of  $x$  in the denominator. What happens then depends on the degrees of the polynomials involved.



**EXAMPLE 8** Numerator and Denominator of 128 / 1564

$$\begin{aligned} \lim_{x \rightarrow \infty} \frac{5x^2 + 8x - 3}{3x^2 + 2} &= \lim_{x \rightarrow \infty} \frac{5}{3} \\ &= \frac{5 + 0 - 0}{3 + 0} = \frac{5}{3} \end{aligned} \quad \text{See Fig. 2.33}$$

**EXAMPLE 9** Degree of Numerator Less Than Degree of Denominator

$$\begin{aligned} \lim_{x \rightarrow \infty} \frac{11x + 2}{2x^3 - 1} &= \lim_{x \rightarrow \infty} \frac{(11/x^2) + (2/x^3)}{2 - (1/x^3)} && \text{Divide numerator and denominator by } x^3 \\ &= \frac{0 + 0}{2 - 0} = 0 && \text{See Fig. 2.34} \end{aligned}$$

We give an example of the case when the degree of the numerator is greater than the degree of the denominator in the next section (Example 8, Section 2.5).

**Horizontal Asymptotes**

If the distance between the graph of a function and some fixed line approaches zero as a point on the graph moves increasingly far from the origin, we say that the graph approaches the line asymptotically and that the line is an *asymptote* of the graph.

Looking at  $f(x) = 1/x$  (See Figure 2.31), we observe that the  $x$ -axis is an asymptote of the curve on the right because

$$\lim_{x \rightarrow \infty} \frac{1}{x} = 0$$

and on the left because

$$\lim_{x \rightarrow -\infty} \frac{1}{x} = 0.$$

We say that the  $x$ -axis is a *horizontal asymptote* of the graph of  $f(x) = 1/x$ .

**DEFINITION** Horizontal Asymptote

A line  $y = h$  is a **horizontal asymptote** of the graph of a function  $y = f(x)$  if either

$$\lim_{x \rightarrow \infty} f(x) = h \quad \text{or} \quad \lim_{x \rightarrow -\infty} f(x) = h.$$

The curve

$$f(x) = \frac{5x^2 + 8x - 3}{3x^2 + 2}$$

sketched in Figure 2.33 (Example 8) has the line  $y = 5/3$  as a horizontal asymptote on both the right and the left because

$$\lim_{x \rightarrow \infty} f(x) = \frac{5}{3} \quad \text{and} \quad \lim_{x \rightarrow -\infty} f(x) = \frac{5}{3}.$$



**EXAMPLE 10** Substituting a New Variable

Find  $\lim_{x \rightarrow \infty} \sin(1/x)$ .

**Solution** We introduce the new variable  $t = 1/x$ . From Example 6, we know that  $t \rightarrow 0^+$  as  $x \rightarrow \infty$  (see Figure 2.31). Therefore,

$$\lim_{x \rightarrow \infty} \sin \frac{1}{x} = \lim_{t \rightarrow 0^+} \sin t = 0.$$

**The Sandwich Theorem Revisited**

The Sandwich Theorem also holds for limits as  $x \rightarrow \pm\infty$ .

**EXAMPLE 11** A Curve May Cross Its Horizontal Asymptote

Using the Sandwich Theorem, find the horizontal asymptote of the curve

$$y = 2 + \frac{\sin x}{x}.$$


**Solution** We are interested in the behavior as  $x \rightarrow \pm\infty$ . Since

$$0 \leq \left| \frac{\sin x}{x} \right| \leq \left| \frac{1}{x} \right|$$

and  $\lim_{x \rightarrow \pm\infty} |1/x| = 0$ , we have  $\lim_{x \rightarrow \pm\infty} (\sin x)/x = 0$  by the Sandwich Theorem. Hence,

$$\lim_{x \rightarrow \pm\infty} \left( 2 + \frac{\sin x}{x} \right) = 2 + 0 = 2,$$

and the line  $y = 2$  is a horizontal asymptote of the curve on both left and right (Figure 2.35).




**FIGURE 2.35** A curve may cross one of its asymptotes infinitely often (Example 11).



**Finding Limits Graphically**

1. Which of the following statements about the function  $y = f(x)$  graphed here are true, and which are false?



- a.  $\lim_{x \rightarrow -1} f(x) = 1$
- b.  $\lim_{x \rightarrow -1} f(x) = 1$
- c.  $\lim_{x \rightarrow -1} f(x)$  exists
- d.  $\lim_{x \rightarrow -1} f(x) = 1$
- e.  $\lim_{x \rightarrow -1} f(x) = 0$
- f.  $\lim_{x \rightarrow -1} f(x)$  does not exist
- g.  $\lim_{x \rightarrow 1} f(x) = 0$
- h.  $\lim_{x \rightarrow 1} f(x) = 0$
- i.  $\lim_{x \rightarrow 1} f(x) = 1$
- j.  $\lim_{x \rightarrow 1} f(x) = 2$
- k.  $\lim_{x \rightarrow 1} f(x)$  does not exist
- l.  $\lim_{x \rightarrow 1} f(x) = 0$

Copyright © 2005 Pearson Education, Inc., publishing as Pearson Addison-Wesley

112 Chapter 2: Limits and Continuity



2. Which of the following statements about the function  $y = f(x)$  graphed here are true, and which are false?




- a.  $\lim_{x \rightarrow 1} f(x) = 1$
- b.  $\lim_{x \rightarrow 1} f(x) = 2$
- c.  $\lim_{x \rightarrow 1} f(x) = 1$
- d.  $\lim_{x \rightarrow 1} f(x) = \lim_{x \rightarrow 1} f(x)$
- e.  $\lim_{x \rightarrow 1} f(x)$  exists at every  $x$  in the open interval  $(-1, 3)$
- f.  $\lim_{x \rightarrow 1} f(x)$  exists at every  $x$  in the open interval  $(-1, 3)$
- g.  $\lim_{x \rightarrow 1} f(x) = 0$
- h.  $\lim_{x \rightarrow 1} f(x)$  does not exist

3. Let  $f(x) = \begin{cases} 3 - x, & x < 2 \\ \frac{1}{2} + x, & x > 2 \end{cases}$



- a. Find  $\lim_{x \rightarrow 2} f(x)$  and  $\lim_{x \rightarrow 2} f(x)$
- b. Does  $\lim_{x \rightarrow 2} f(x)$  exist? If so, what is it? If not, why not?
- c. Find  $\lim_{x \rightarrow 2} f(x)$  and  $\lim_{x \rightarrow 2} f(x)$
- d. Does  $\lim_{x \rightarrow 2} f(x)$  exist? If so, what is it? If not, why not?

4. Let  $f(x) = \begin{cases} 3 - x, & x < 2 \\ 2, & x = 2 \\ \frac{1}{2} + x, & x > 2 \end{cases}$




- a. Find  $\lim_{x \rightarrow 2} f(x)$  and  $\lim_{x \rightarrow 2} f(x)$
- b. Does  $\lim_{x \rightarrow 2} f(x)$  exist? If so, what is it? If not, why not?
- c. Find  $\lim_{x \rightarrow 2} f(x)$  and  $\lim_{x \rightarrow 2} f(x)$
- d. Does  $\lim_{x \rightarrow 2} f(x)$  exist? If so, what is it? If not, why not?

Copyright © 2005 Pearson Education, Inc., publishing as Pearson Addison-Wesley

- a. Find  $\lim_{x \rightarrow 0} f(x)$ ,  $\lim_{x \rightarrow 0} f(x)$ , and  $f(0)$
- b. Does  $\lim_{x \rightarrow 0} f(x)$  exist? If so, what is it? If not, why not?
- c. Find  $\lim_{x \rightarrow 0} f'$
- d. Does  $\lim_{x \rightarrow 0} f'$


131-132 / 1564

5. Let  $f(x) = \begin{cases} 0, & x = 0 \\ \sin \frac{1}{x}, & x \neq 0 \end{cases}$



- a. Does  $\lim_{x \rightarrow 0} f(x)$  exist? If so, what is it? If not, why not?
- b. Does  $\lim_{x \rightarrow 0} f(x)$  exist? If so, what is it? If not, why not?
- c. Does  $\lim_{x \rightarrow 0} f(x)$  exist? If so, what is it? If not, why not?

6. Let  $g(x) = \sqrt{x} \sin(1/x)$



- a. Does  $\lim_{x \rightarrow 0} g(x)$  exist? If so, what is it? If not, why not?
- b. Does  $\lim_{x \rightarrow 0} g(x)$  exist? If so, what is it? If not, why not?
- c. Does  $\lim_{x \rightarrow 0} g(x)$  exist? If so, what is it? If not, why not?

Copyright © 2005 Pearson Education, Inc., publishing as Pearson Addison-Wesley



- 7. a. Graph  $f(x) = \begin{cases} x^2, & x \neq 1 \\ 0, & x = 1 \end{cases}$
- b. Find  $\lim_{x \rightarrow 1} f(x)$  and  $\lim_{x \rightarrow 1} f(x)$
- c. Does  $\lim_{x \rightarrow 1} f(x)$  exist? If so, what is it? If not, why not?
- 8. a. Graph  $f(x) = \begin{cases} 1 - x^2, & x \neq 1 \\ 2, & x = 1 \end{cases}$
- b. Find  $\lim_{x \rightarrow 1} f(x)$  and  $\lim_{x \rightarrow 1} f(x)$
- c. Does  $\lim_{x \rightarrow 1} f(x)$  exist? If so, what is it? If not, why not?

Graph the functions in Exercises 9 and 10. Then answer these questions.

- a. What are the domain and range of  $f$ ?
- b. At what points  $x$ , if any, does  $\lim_{x \rightarrow c} f(x)$  exist?
- c. At what points does only the left-hand limit exist?
- d. At what points does only the right-hand limit exist?



- 9.  $f(x) = \begin{cases} \sqrt{1-x^2}, & 0 \leq x < 1 \\ 1, & 1 \leq x < 2 \\ 2, & x = 2 \end{cases}$
- 10.  $f(x) = \begin{cases} x - 1, & x < 0, \text{ or } 0 < x < 1 \\ 1, & x = 0 \\ 0, & x < -1, \text{ or } x > 1 \end{cases}$

**Finding One-Sided Limits Algebraically**

Find the limits in Exercises 11–18.



- 11.  $\lim_{x \rightarrow 1} \frac{x^2 - 1}{x + 1}$
- 12.  $\lim_{x \rightarrow 1} \frac{x^2 - 1}{x - 2}$
- 13.  $\lim_{x \rightarrow 2} \left( \frac{x}{x+1} \right) \left( \frac{x^2 - 4}{x - 2} \right)$
- 14.  $\lim_{x \rightarrow 1} \left( \frac{1}{x-1} \right) \left( \frac{x^2 - 1}{x+1} \right) \left( \frac{1}{x-1} \right)$
- 15.  $\lim_{x \rightarrow 0} \frac{\sqrt{x^2 + 4x + 9} - \sqrt{9}}{x}$
- 16.  $\lim_{x \rightarrow 0} \frac{\sqrt{x^2 + 5x + 16} - 4}{x}$
- 17. a.  $\lim_{x \rightarrow 2} (x + 3) \left( \frac{x + 2}{x - 2} \right)$
- b.  $\lim_{x \rightarrow 2} (x + 3) \left( \frac{x + 2}{x - 2} \right)$
- 18. a.  $\lim_{x \rightarrow 1} \frac{\sqrt{2x(x-1)}}{(x-1)}$
- b.  $\lim_{x \rightarrow 1} \frac{\sqrt{2x(x-1)}}{(x-1)}$

Use the graph of the greatest integer function  $y = \lfloor x \rfloor$  (sometimes written  $y = \text{int } x$ ) in Figure 1.11 in Section 1.1, to help you find the limits in Exercises 19 and 20.

- 19. a.  $\lim_{x \rightarrow 1} \frac{\lfloor x \rfloor}{x}$
- b.  $\lim_{x \rightarrow 1} \frac{\lfloor x \rfloor}{x}$
- 20. a.  $\lim_{x \rightarrow 1} (\lfloor x \rfloor)$
- b.  $\lim_{x \rightarrow 1} (\lfloor x \rfloor)$

Copyright © 2005 Pearson Education, Inc., publishing as Pearson Addison-Wesley

2.4 One-Sided Limits and Limits at Infinity 113

**Using  $\lim_{\theta \rightarrow 0} \frac{\sin \theta}{\theta} = 1$**

Find the limits in Exercises 21–36.

- 21.  $\lim_{x \rightarrow 0} \frac{\sin \sqrt{x}}{\sqrt{x}}$
- 22.  $\lim_{x \rightarrow 0} \frac{\sin 3x}{3x}$  (a constant)
- 23.  $\lim_{x \rightarrow 0} \frac{\sin 5x}{5x}$
- 24.  $\lim_{x \rightarrow 0} \frac{\sin 7x}{7x}$
- 25.  $\lim_{x \rightarrow 0} \frac{\sin 2x}{2x}$
- 26.  $\lim_{x \rightarrow 0} \frac{\sin x}{x}$
- 27.  $\lim_{x \rightarrow 0} \frac{x \cos 2x}{\sin 3x}$
- 28.  $\lim_{x \rightarrow 0} \frac{\sqrt{x} \cos x}{\sin 2x}$
- 29.  $\lim_{x \rightarrow 0} \frac{1 - \cos x}{x \sin x}$
- 30.  $\lim_{x \rightarrow 0} \frac{x^2 - x + \sin x}{2x}$
- 31.  $\lim_{x \rightarrow 0} \frac{\sin(1 - \cos x)}{1 - \cos x}$
- 32.  $\lim_{x \rightarrow 0} \frac{\sin(\sin x)}{\sin x}$
- 33.  $\lim_{x \rightarrow 0} \frac{\sin \theta}{\theta}$
- 34.  $\lim_{x \rightarrow 0} \frac{\sin 9x}{\sin 4x}$
- 35.  $\lim_{x \rightarrow 0} \frac{\sin 7x}{\sin 6x}$
- 36.  $\lim_{x \rightarrow 0} \frac{\sin 3x \cos 5x}{x \sin 4x}$

**Calculating Limits as  $x \rightarrow \pm \infty$**

In Exercises 37–42, find the limit of each function (a) as  $x \rightarrow \infty$  and (b) as  $x \rightarrow -\infty$ . (You may wish to visualize your answer with a graphing calculator or computer.)

- 37.  $f(x) = \frac{x}{2} - 3$
- 38.  $f(x) = x - \frac{2}{x^2}$
- 39.  $g(x) = \frac{1}{2 + (1/x)}$
- 40.  $g(x) = \frac{1}{8 - (3/x^2)}$
- 41.  $h(x) = \frac{-5 + (7/x)}{3 - (1/x^2)}$
- 42.  $h(x) = \frac{3 - (2/x)}{4 + (\sqrt{2}/x^2)}$

Find the limits in Exercises 43–46.

- 43.  $\lim_{x \rightarrow \infty} \frac{\sin 2x}{2x}$
- 44.  $\lim_{x \rightarrow \infty} \frac{\cos \theta}{\theta}$
- 45.  $\lim_{x \rightarrow \infty} \frac{2 - x + \sin x}{x + \cos x}$
- 46.  $\lim_{x \rightarrow \infty} \frac{x + \sin x}{2x^2 + 7 - 3 \sin x}$

**Limits of Rational Functions**

In Exercises 47–56, find the limit of each rational function (a) as  $x \rightarrow \infty$  and (b) as  $x \rightarrow -\infty$ .

- 47.  $f(x) = \frac{2x + 1}{3x + 7}$
- 48.  $f(x) = \frac{2x^2 + 7}{x^2 - x + 7}$
- 49.  $f(x) = \frac{x + 1}{x^2 + 3}$
- 50.  $f(x) = \frac{3x + 9}{x^2 - 2}$
- 51.  $h(x) = \frac{1}{x^2 - 3x + 6}$
- 52.  $g(x) = \frac{1}{x^2 - 4x + 1}$

Copyright © 2005 Pearson Education, Inc., publishing as Pearson Addison-Wesley